

## PURITY OF BOUNDARIES OF OPEN COMPLEX VARIETIES

ANDRZEJ WEBER

ABSTRACT. We study the boundary of an open smooth complex algebraic variety  $U$ . We ask when the cohomology of the geometric boundary  $Z = X \setminus U$  in a smooth compactification  $X$  is pure with respect to the mixed Hodge structure. Knowing the dimension of singularity locus of some singular compactification, we give a bound for  $k$  above which the cohomology  $H^k(Z)$  is pure. The main ingredient of the proof is purity of the intersection cohomology sheaf.

### 1. INTRODUCTION

Let  $U$  be a smooth complex algebraic variety which is not compact. We study cohomological properties of  $U$  which are invariant with respect to modifications of the interior of  $U$ . In other words, we investigate cohomological properties of the boundary.

The boundary itself can have at least two meanings. First of all, from the topological point of view, we may treat an open smooth variety as the interior of a compact manifold with boundary. In this case, the boundary would mean an odd-dimensional real manifold. We call it the link at infinity. On the other hand, from the geometric point of view, we may compactify our variety in the category of algebraic varieties. The boundary is then a subvariety of the compactification. In addition, we may require that the compactification is smooth. The condition that the boundary is a normal crossing divisor is irrelevant to us, although it is hidden in the construction of the mixed Hodge structure. The link at infinity can be identified with the link of the boundary in the compactification. Despite the differences, we show that the topological and geometric boundaries have a great deal in common where the mixed Hodge structure is concerned.

To some extent, we try to avoid specific methods of Hodge theory, having in mind possible applications (or rather open questions) to real algebraic geometry, as well as some questions about torsion for cohomology of complex varieties. The sections §2-§4 are valid in that generality. Nevertheless, the results of §5 cannot be generalized and they hold only for rational cohomology of algebraic varieties. The strong functoriality of the weight filtration implies that the lower weight subspaces of the topological and geometric boundaries coincide; see Proposition 3. In similar situations, this phenomenon was already described in [10, Prop. 7.1] and [2, Prop. 5.1].

The proof of the main result of §6 uses even stronger techniques. The purity of the intersection cohomology sheaf [3] imposes some conditions on the link of the geometric boundary. We prove Theorem 11, which can be shortened to the following statement:

**Theorem 1.** *Suppose  $U$  is a complex smooth algebraic variety. Assume that  $U$  admits a singular compactification  $Y$ . Suppose that the singularity of the pair  $(Y, Y \setminus U)$  is of dimension  $s$ . Then, for every smooth compactification  $X$ , the boundary  $X \setminus U$  has pure cohomology  $H^k(X \setminus U)$ , for  $k \geq \dim(U) + s$ .*

---

2000 *Mathematics Subject Classification.* 32S35, 55N33, 14E15, 32S20.

*Key words and phrases.* Mixed Hodge theory of singular varieties, intersection cohomology, resolution of singularities.

Theorem 1, in the case where  $U$  admits a one-point compactification, already appeared in [4, Th. 2.1.11]. A vast generalization was given in [13]. The present version gives a better bound for purity, although the situation considered here is less general.

One can treat Theorem 1 as a contractibility criterion. For a subvariety  $Z \subset X$ : if  $H^k(Z)$  is not pure for  $k \geq \dim(U) + s$ , then the pair  $(X, Z)$  cannot be contracted to a pair  $(Y, W)$  with singularities of dimension less than or equal to  $s$ . Although Theorem 1 resembles the Grauert criterion, it is of a different nature. In the Grauert criterion, the intersection form on  $Z$  depends on the embedding  $Z \subset X$ , whereas, here, the mixed Hodge structure of  $Z$  does not. Needless to say, our criterion is not sufficient for the existence of a contraction.

In contrast to our previous paper [13], we try to present the subject in a form as elementary as possible. We have avoided using mixed the Hodge modules of [11], taking for granted that the cohomology with coefficients in a complex of sheaves of geometric origin has a natural mixed Hodge structure. By “geometric origin”, we mean “obtained by the standard sheaf-theoretic operations”. We assume that the varieties are defined over the real numbers or over the complex numbers, and we use classical topology. In our arguments, we will apply resolution of singularities, although parts of the results depend only on the formal properties of mixed Hodge modules or Weil sheaves.

## 2. TOPOLOGICAL BOUNDARY: THE LINK AT INFINITY

Let us begin with a description of some invariants of open manifolds which can be defined using just topology and basic properties of resolutions of singularities. Working with  $\mathbb{Z}/2$ -cohomology, we can apply our construction also for real algebraic manifolds. For complex manifolds, we can use any coefficients, not necessarily  $\mathbb{Q}$ .

The first invariant we propose to consider is the cohomology of the link at infinity:

$$H^*(L_\infty U) := \varinjlim_{K \subset U} H^*(U \setminus K),$$

where  $K$  runs through compact sets contained in  $U$ . The group  $H^*(L_\infty U)$  is exactly the cohomology of the link  $L_Z$ , which is the link of the boundary set  $Z = X \setminus U$ , where  $X$  is a compactification of  $U$ . (For various approaches to the link of a subvariety, see [5].) This cohomology group is of finite dimension. It can be expressed in terms of sheaf operations on  $X$ :

$$H^*(L_\infty U) = H^*(L_Z) = H^*(Z; i^* Rj_* \mathbb{Q}_U),$$

where  $j : U \hookrightarrow X$  and  $i : X \setminus U \hookrightarrow X$  are the inclusions.

## 3. GEOMETRIC BOUNDARY: IMAGE OF BOUNDARY CYCLES

Another invariant which we consider is the image of the boundary cocycles

$$IB^*(U) = \text{im}(H^*(Z) \rightarrow H_c^{*+1}(U)) = \ker(H_c^{*+1}(U) \rightarrow H^{*+1}(X)),$$

where  $X$  is a **smooth** compactification of  $U$  and  $Z = X \setminus U$ . The maps come from the long exact sequence of the pair  $(X, Z)$ .

To show the independence of  $X$ , we start with a purely topological lemma.

**Lemma 2.** *Suppose we have a map of real smooth oriented closed manifolds*

$$f : X_1 \rightarrow X_2,$$

*which is an isomorphism of some open subsets*

$$f|_{U_1} : U_1 = f^{-1}(U_2) \xrightarrow{\cong} U_2.$$

Denote by

$$IB_i^* = \ker(H_c^{*+1}(U_i) \rightarrow H^{*+1}(X_i))$$

the kernels of the natural maps for  $i = 1, 2$ . Then  $f$  induces an isomorphism

$$f^* : IB_2^* \rightarrow IB_1^* .$$

**Proof.** The map  $f^* : H^*(X_2) \rightarrow H^*(X_1)$  is injective, since it is a map of degree one of compact manifolds. The map  $f$  induces the transformation

$$\begin{array}{ccccc} IB_2^k & \hookrightarrow & H_c^{k+1}(U_2) & \rightarrow & H^{k+1}(X_2) \\ & & \simeq \downarrow & & \text{mono} \downarrow \\ IB_1^k & \hookrightarrow & H_c^{k+1}(U_1) & \rightarrow & H^{k+1}(X_1) . \end{array}$$

It follows that  $IB_2^k \rightarrow IB_1^k$  is an isomorphism.  $\square$

To prove the independence of  $IB^*(U)$  on the compactification, simply note that any two smooth compactifications are dominated by a third one.

#### 4. BASIC EXACT SEQUENCES

We will need three exact sequences to relate the described invariants. These exact sequences may be constructed topologically, but it is important to know that they come from distinguished triangles in the derived category of sheaves. It will follow that, for complex varieties, the maps of the described exact sequences preserve the mixed Hodge structure.

We start with the sequence relating the cohomology of  $U$  and the cohomology of its link at infinity. Let  $X$  be any compactification and  $Z = X \setminus U$ . It is possible to find a neighbourhood  $N$  of  $Z$  which retracts to  $Z$  and such that the boundary  $\partial N$  is homeomorphic to the link of  $Z$ . Considering the pair  $(X \setminus N, \partial N)$ , we arrive at the long exact sequence

$$(1) \quad \dots \rightarrow H^k(U) \rightarrow H^k(L_Z) \xrightarrow{\delta} H_c^{k+1}(U) \rightarrow H^{k+1}(U) \rightarrow \dots .$$

This exact sequence may, in fact, be obtained from the fundamental distinguished triangle (in the category of mixed Hodge modules on  $X$ )

$$(2) \quad \begin{array}{ccc} i_* i^! G & \longrightarrow & G \\ [+1] \nearrow & & \nwarrow \\ & Rj_* j^* G & \end{array}$$

where  $G = j_! \mathbb{Q}_U$ . By duality, we obtain the triangle

$$\begin{array}{ccc} i_! i^* Rj_* \mathbb{Q}_U & \longleftarrow & Rj_* \mathbb{Q}_U \\ [+1] \searrow & & \nearrow \\ & j_! \mathbb{Q}_U & \end{array}$$

since  $j_! j^! Rj_* \mathbb{Q}_U \simeq j_! \mathbb{Q}_U$ . Applying cohomology, we obtain the sequence (1).

We also need an exact sequence relating  $H^k(Z)$  and  $H^k(L_Z)$ . Topologically, we have a deformation retraction from the closure  $\bar{N} \rightarrow Z$ . The exact sequence for the manifold with boundary  $(\bar{N}, \partial N)$ ,

$$\dots \rightarrow H^k(\bar{N}) \rightarrow H^k(\partial N) \rightarrow H^{k+1}(\bar{N}, \partial N) \rightarrow H^{k+1}(\bar{N}) \rightarrow \dots ,$$

becomes

$$(3) \quad \dots \rightarrow H^k(Z) \rightarrow H^k(L_Z) \rightarrow H^{k+1}(X, U) \rightarrow H^{k+1}(Z) \rightarrow \dots .$$

The sheaf theoretic definition is given below. Let us restrict the triangle (2) with  $G = \mathbb{Q}_X$  to  $Z$ . We have  $i^*i_*i^!\mathbb{Q}_X = i^!\mathbb{Q}_X$ , and we obtain the triangle

$$\begin{array}{ccc} i^!\mathbb{Q}_X & \longrightarrow & \mathbb{Q}_Z \\ [+1] \swarrow & & \swarrow \\ & i^*Rj_*\mathbb{Q}_U & \end{array}$$

The associated sequence of cohomology is just (3). It plays a fundamental role in our further consideration.

Of course, the third exact sequence that we use is the sequence of the pair  $(X, Z)$

$$(4) \quad \cdots \rightarrow H^k(X) \rightarrow H^k(Z) \rightarrow H_c^{k+1}(U) \rightarrow H^{k+1}(X) \rightarrow \cdots .$$

To relate the groups  $H^*(L_\infty(U)) = H^*(L_Z)$  and  $IB^*(U) = \text{im}(H^k(Z) \rightarrow H_c^{k+1}(U))$ , we apply the map of exact sequences from (4) to (1) induced by the inclusion

$$(X \setminus N, \partial N) \subset (X, N) \xrightarrow{htp} (X, Z) .$$

We obtain the commutative diagram

$$\begin{array}{ccc} H^k(Z) & \rightarrow & H_c^{k+1}(U) \\ \downarrow & & \parallel \\ H^k(L_Z) & \rightarrow & H_c^{k+1}(U) \end{array} .$$

We see that

$$IB^k(U) = \text{im}(H^k(Z) \rightarrow H_c^{k+1}(U)) \subset \text{im}(H^k(L_\infty(U)) \rightarrow H_c^{k+1}(U)) .$$

In general, this inclusion is proper.

## 5. MIXED HODGE STRUCTURE

From now on, we consider only complex algebraic varieties and rational cohomology.

The invariants  $H^k(L_\infty U)$  and  $IB^k(U)$  are equipped with mixed Hodge structures. The first one is given by the sheaf-theoretic description:

$$H^*(L_\infty U) = H^*(Z; i^*Rj_*\mathbb{Q}_U) .$$

The second one,  $IB^*(U)$ , has a structure induced from  $H^*(Z)$ . In the situation of Lemma 2, the map  $IB_2^k \rightarrow IB_1^k$  preserves the quotient mixed Hodge structures and since it is an isomorphism of vector spaces it must be also an isomorphism of all weight subspaces. In fact, by the definition of the mixed Hodge structure, we have

$$IB^k(U) = W_k H_c^{k+1}(U) .$$

For us, the most interesting part is the weight subspace  $W_{k-1}$ . Using basic properties of the mixed Hodge structure, we will give three descriptions of that weight space.

**Proposition 3.** *Let  $X$  be a smooth compactification of  $U$  and  $Z = X \setminus U$ . Then the following groups are isomorphic:*

- (1)  $W_{k-1}H^k(L_\infty U)$ ,
- (2)  $W_{k-1}H_c^{k+1}(U)$ ,
- (3)  $W_{k-1}H^k(Z)$ .

Note that, in the statement of the theorem, we do not assume that  $Z$  is a smooth divisor with a normal crossings. As a corollary to Proposition 3, we have

**Corollary 4.** *Let  $X$  be a smooth compactification of  $U$  and  $Z = X \setminus U$ . The cohomology  $H^k(Z)$  is pure, of weight  $k$ , if and only if  $H^k(L_Z)$  is of weight  $\geq k$ .*

Also we note (compare [10, Prop. 7.1]):

**Corollary 5.** *The impure part of cohomology of the boundary set  $W_{k-1}H^k(Z)$  does not depend on the smooth compactification.*

**Remark 6.** Note that the group  $W_{k-1}H^k(Z)$  is a topological invariant of  $Z$ , since, by [12], it is the kernel of the canonical map to intersection cohomology  $H^k(Z) \rightarrow IH^k(Z)$ . Also, by the construction of the mixed Hodge structure, we have

$$W_{k-1}H^k(Z) = \ker(g^* : H^k(Z) \rightarrow H^k(\tilde{Z})),$$

where  $g : \tilde{Z} \rightarrow Z$  is any dominating proper map from a smooth variety, possibly of bigger dimension.

The entire cohomology of the boundary of a smooth compactification is not an invariant of  $U$ . Of course, when we blow up something at the boundary, the cohomology is modified; nevertheless, the lower parts of the weight filtration remain unchanged.

**Remark 7.** With the help of the Decomposition Theorem of [3], we have better insight into what happens with the cohomology of the boundary. Let  $f$  be a map of pairs  $(X_1, Z_1) \rightarrow (X_2, Z_2)$  which is an isomorphism outside  $Z_1$ . The push-forward of the constant sheaf on  $X_1$  decomposes:

$$Rf_*\mathbb{Q}_{X_1} \simeq \mathbb{Q}_{X_2} \oplus \bigoplus_{\alpha} IC(V_{\alpha}; L_{\alpha}).$$

The supports of the intersection sheaves  $IC(V_{\alpha}; L_{\alpha})$  are contained in  $Z_2$ ; therefore

$$H^*(Z_1) = H^*(Z_2; (Rf_*\mathbb{Q}_{X_1})|_{Z_2}) \simeq H^*(Z_2) \oplus \bigoplus_{\alpha} IH^*(V_{\alpha}; L_{\alpha}).$$

Again, we see that the difference between  $H^*(Z_1)$  and  $H^*(Z_2)$  is pure, since  $\bigoplus_{\alpha} IH^*(V_{\alpha}; L_{\alpha})$  is a summand of  $H^*(X_1)$ .

**Remark 8.** Using another powerful tool, namely the Weak Factorization Theorem [1], we can trace how the cohomology of the boundary may change. Each time, when we blow up a smooth center  $S$  contained in the boundary, the pure summand  $\operatorname{coker}(H^*(S) \rightarrow H^*(\mathbb{P}N_{S/X}))$  contributes to the cohomology of the blown-up boundary. Here,  $H^*(\mathbb{P}N_{S/X})$  is the projectivization of the normal bundle of  $S$  in  $X$ .

The proof of Proposition 3 is divided into Lemmas 9 and 10.

**Lemma 9.** *We have*

$$W_{k-1}H^k(Z) \simeq W_{k-1}H_c^{k+1}(U).$$

**Proof.** We recall that  $H^k(X)$  is of weight  $k$  and  $H^{k+1}(X)$  is of weight  $k + 1$ . Therefore, the long exact sequence

$$\cdots \rightarrow H^k(X) \rightarrow H^k(Z) \xrightarrow{\delta} H_c^{k+1}(U) \rightarrow H^{k+1}(X) \rightarrow \cdots$$

induces an isomorphism of graded pieces for  $\ell < k$

$$Gr_{\ell}^W H^k(Z) \simeq Gr_{\ell}^W H_c^{k+1}(U).$$

It follows that the boundary map  $W_{k-1}H^k(Z) \rightarrow W_{k-1}H_c^{k+1}(U)$  is an isomorphism.  $\square$

**Lemma 10.** *We have*

$$W_{k-1}H^k(L_Z) \simeq W_{k-1}H_c^{k+1}(U).$$

**Proof.** We consider the long exact sequence (1). Since  $U$  is smooth,  $W_{k-1}H^k(U) = 0$ . Therefore, for  $\ell < k$

$$Gr_\ell^W H^k(L_Z) \simeq Gr_\ell^W H_c^{k+1}(U).$$

Again, the boundary map  $W_{k-1}H^k(L_Z) \rightarrow W_{k-1}H_c^{k+1}(U)$  is an isomorphism.  $\square$

6. SINGULAR VERSUS SMOOTH COMPACTIFICATIONS

Let  $W \subset Y$  be a pair of varieties. Assume that  $Y \setminus W$  is smooth. By the “singularity of the pair”, we mean the set of points at which  $W$  in  $Y$  analytically does not look like a submanifold (of any dimension) in a manifold. The singularity set consists of points at which  $W$  or  $Y$  is singular. Below, we give the exact statement of our main result.

**Theorem 11.** *Let  $U$  be a smooth variety. Suppose that  $U$  admits a compactification  $Y$  and let  $W = Y \setminus U$  be the boundary set. Denote by  $s$  the dimension of the singularity of the pair  $(Y, W)$ . Let  $X$  be a smooth compactification of  $U$  and  $Z = X \setminus U$ . For  $k \geq \dim(U) + \dim(W)$ , we have:*

- i) the cohomology of the link  $H^k(L_Z)$  is of weight  $\geq k + 1$ ,*
- ii) the restriction map  $H^k(Z) \rightarrow H^k(L_Z)$  vanishes.*

For  $k \geq \dim(U) + s$  we have:

- iii) the cohomology of the boundary  $H^k(Z)$  is pure, of weight  $k$ , that is*

$$W_{k-1}H^k(Z) = 0$$

- iv) the cohomology of the link  $H^k(L_Z)$  is of weight  $\geq k$ .*

Note that, by Proposition 3, claim iv) does not depend on the choice of the smooth compactification  $X$ .

Let  $n = \dim(U)$ . By Poincaré duality, we have

$$\begin{aligned} H^k(Z)^* &= H^{2n-k}(X, U)(n), \\ H^k(L_Z)^* &= H^{2n-1-k}(L_Z)(n), \end{aligned}$$

where  $(n)$  denotes the Tate twist shifting the weights by  $2n$ . The dual version of Theorem 11 is the following:

**Theorem 12.** *With the assumptions of Theorem 11:*

For  $k \leq \dim(U) - \dim(W)$ , we have

- i') the cohomology of the link  $H^{k-1}(L_Z)$  is of weight  $\leq k - 1$ ,*
- ii') the boundary map  $H^{k-1}(L_Z) \rightarrow H^k(X, U)$  vanishes.*

For  $k \leq \dim(U) - s$ , we have

- iii') the cohomology  $H^k(X, U)$  is pure, of weight  $k$ , that is*

$$W_k H^k(X, U) = H^k(X, U)$$

- iv') the cohomology of the link  $H^{k-1}(L_Z)$  is of weight  $\leq k$ .*

To distinguish two copies of  $U$  in  $X$  and in  $Y$ , we will use the letter  $V$  for the copy of  $U$  in  $Y$ . The identification map  $U \rightarrow V$  is denoted by  $f$ :

$$\begin{array}{ccccc} Z = X \setminus U & \subset & X = \bar{U} & \supset & U \\ & & & \simeq & \downarrow f \\ W = Y \setminus V & \subset & Y = \bar{V} & \supset & V \end{array}$$

**Remark 13.** In our setup, we can apply completion and resolution of singularities. Therefore,  $X$  can be replaced by a dominating smooth variety for which the map  $f$  extends to the boundary.

Some information about the weights of the cohomology of the link and the boundary can be deduced when we have a proper map  $f : U \rightarrow V$  and a compactification of  $V$ . A statement which generalizes i) and ii) in terms of the *defect of semismallness* (introduced in [4]) is formulated in [13]. The direct generalization of iii) and iv) would involve precise information about the singularities of the perverse cohomology sheaves  ${}^p\mathcal{H}^k Rf_*\mathbb{Q}_U$ .

Theorem 11 can be localized around a topological component of  $X \setminus U$ . Precisely, consider the set of ends, i.e.,  $U_\infty = \pi_0(X \setminus U)$ . This set does not depend on the choice of  $X$ , provided that  $X$  is normal. A map of algebraic varieties which is proper induces a map of their ends. To deduce purity of the cohomology of a part of the boundary of  $U$ , it is enough to have information about a singular completion of the corresponding end.

## 7. PROOFS

Before the proof of Theorem 11, let us recall the key property of the link of a subvariety.

**Theorem 14** ([5]). *Let  $Y$  be a variety and let  $W$  be a compact subvariety. Let us assume that  $Y \setminus W$  is smooth. Then,  $H^k(L_W)$  is of weight  $\leq k$ , for degrees  $k < \dim(Y) - \dim(W)$ .*

Theorem 14 immediately follows from the purity of the intersection sheaf [6, 3], since the stalk cohomology  $\mathcal{H}^k(IC_Y)$  is isomorphic to  $\mathcal{H}^k(Rj_*\mathbb{Q}_V)$  for  $k < \dim(Y) - \dim(W)$  and  $H^*(L_W) = H^*(W; (Rj_*\mathbb{Q}_V)|_W)$ .

**Remark 15.** In [2, §6], the Decomposition Theorem of [3] was used to give estimates for the dimension of intersection cohomology of the link by means of resolution. But it seems that the purity of the intersection sheaf was not used directly.

**Proof of (12.i'-ii').**

By Remark 13, we assume that the map  $f$  extends to  $X$ . The extended map (denoted by the same letter) induces a map of sheaves  $i'^*Rj'_*\mathbb{Q}_V \rightarrow Rf_*i^*Rj_*\mathbb{Q}_U$ . Therefore, the mixed Hodge structures of the isomorphic groups  $H^*(L_W)$  and  $H^*(L_Z)$  coincide. By Theorem 14 and the assumption on the dimension of  $W$ , the cohomology  $H^{k-1}(L_W)$  is of weight  $\leq k - 1$ . Claim 12.ii' follows from the long exact sequence (3): the boundary map

$$H^{k-1}(L_Z) \rightarrow H^k(X, U)$$

vanishes because the first term is of weight  $\leq k - 1$  and the second term is of weight  $\geq k$ .

**Proof of (11.i-ii)** follows by duality.

**Proof of (11.iii-iv)** If  $W = \text{Sing}(Y)$ , then  $s = \dim(W)$ , and statement i) is even stronger than iii). Also by Proposition 3 we have  $W_{k-1}H^k(Z) = W_{k-1}H^k(L_Z) \subset W_kH^k(L_Z) = 0$ , therefore iv) follows.

Suppose now that  $\text{Sing}(Y) \subsetneq W$ . We may assume that  $f$  extends to a map  $X \rightarrow Y$  and also we may assume that the map  $f$  is a resolution of singularities of the pair  $(Y, W)$ . Let  $\widetilde{W} \subset X$  be the proper transform of  $W$ . Denote by  $E \subset X$  the exceptional set of  $f$  and let  $F = E \cap \widetilde{W}$ . Consider the Mayer-Vietoris exact sequence for  $Z = E \cup \widetilde{W}$ :

$$\dots \rightarrow H^{k-1}(E) \oplus H^{k-1}(\widetilde{W}) \xrightarrow{\alpha} H^{k-1}(F) \xrightarrow{\delta} H^k(Z) \rightarrow H^k(E) \oplus H^k(\widetilde{W}) \rightarrow \dots$$

By (11.i), applied to the map  $(X, E) \rightarrow (Y, f(E))$ , the cohomology of the link  $H^k(L_E)$  is of weight  $\geq k + 1$  for  $k \geq \dim(X) + s$ . Hence, by Proposition 3, the cohomology  $H^k(E)$  is pure for

$k \geq \dim(X) + s$ . Of course,  $H^k(\widetilde{W})$  is pure since we assume that  $\widetilde{W}$  is smooth. To prove the purity of  $H^k(Z)$ , it remains to show that the map  $\delta$  of the Mayer-Vietoris sequence is trivial.

By (11.ii), applied to  $F \subset \widetilde{W}$ , the map

$$H^{k-1}(F) \rightarrow H^{k-1}(L_F)$$

vanishes for  $k - 1 \geq \dim(W) + s$ . By the exact sequence (3) of that pair, the restriction map

$$H^{k-1}(\widetilde{W}, \widetilde{W} \setminus F) \rightarrow H^{k-1}(F)$$

is surjective. The above map factors through  $H^{k-1}(\widetilde{W})$ ; therefore, the map

$$H^{k-1}(\widetilde{W}) \rightarrow H^{k-1}(F)$$

is surjective. It follows that the restriction map  $\alpha$  is surjective and the boundary map  $\delta$  is trivial for  $k \geq \dim(X) + s \geq \dim(W) + s + 1$ . This completes the proof.

**Proof of (12.iii'-iv')** follows by duality.  $\square$

**Remark 16.** If the singularity set is empty, then  $s = -\infty$  by convention. Claims (11.i-ii) hold in all degrees for trivial reasons.

The special case when  $W$  is a point (an isolated singularity resolution) was studied from the very beginning of the theory. In that case, both maps  $H^{n-1}(L_Z) \rightarrow H^n(X, U)$  and  $H^n(X) \rightarrow H^n(L_Z)$  are trivial. The map  $H^n(X, U) \rightarrow H^n(Z)$  is an isomorphism. After the identification  $H^n(X, U) = H^n(Z)^*$ , we obtain a nondegenerate intersection form, which was studied for example in [7].

## 8. QUESTIONS ABOUT REAL ALGEBRAIC VARIETIES

Hodge theory for real algebraic varieties and  $\mathbb{Z}/2$  coefficients is not available. The approach of [8, 9] does not lead to a strongly functorial weight filtration. Nevertheless, one defines impure cohomology of a singular compact variety  $X$ : it is the kernel of  $H^*(X) \rightarrow H^*(\tilde{X})$ , where  $\tilde{X}$  is any resolution. We say that the cohomology of a real variety is pure if the kernel  $H^*(X) \rightarrow H^*(\tilde{X})$  is trivial. The definition does not depend on  $\tilde{X}$ . Now one can ask the question about a generalization of Theorem 11:

**Question 17.** *With the assumption of Theorem 3 for real algebraic varieties: What properties of  $(Y, W)$  would guarantee purity of  $H^*(Z)$  in some range of degrees?*

The dimension of the singularity set is far too weak of an invariant. It is well-known that any real algebraic set can be contracted to a point.

## REFERENCES

- [1] Abramovich, D.; Karu, K.; Matsuki, K.; Włodarczyk, J.: *Torification and factorization of birational maps*. J. Am. Math. Soc. 15, No.3, 531-572 (2002) DOI: [10.1090/S0894-0347-02-00396-X](https://doi.org/10.1090/S0894-0347-02-00396-X)
- [2] Arapura D.; Bakhtary, P.; Włodarczyk J.: *Weights on cohomology, invariants of singularities, and dual complexes*. [arXiv:1102.4370](https://arxiv.org/abs/1102.4370)
- [3] Beilinson, A. A.; Bernstein, J.; Deligne, P.: *Faisceaux pervers*. Analysis and topology on singular spaces, I (Luminy, 1981), 5-171, Astérisque, 100, Soc. Math. France, Paris, 1982.
- [4] de Cataldo, M. A.; Migliorini, L.: *The Hodge theory of algebraic maps*, Ann. Sci. Éc. Norm. Supér. (4) 38 (2005), no. 5, 693-750.
- [5] Durfee, A. H.; Saito, M.: *Mixed Hodge structures on the intersection cohomology of links*, Compos. Math. 76 (1990), no.1/2, 49-67.
- [6] Gabber, O.: Preprint IHES 1980

- [7] Goresky, M.; MacPherson, R.: *On the topology of complex algebraic maps*. Algebraic geometry (La Rábida, 1981), 119-129, Lecture Notes in Math., 961, Springer, Berlin, 1982.
- [8] McCrory, C.; Parusinski, A.: *The weight filtration for real algebraic varieties II: Classical homology*. Friedman, Greg (ed.) et al., Topology of stratified spaces. Cambridge University Press. Mathematical Sciences (2011), 121-160
- [9] McCrory, C.; Parusinski, A.: *The weight filtration for real algebraic varieties II: Classical homology*. [arXiv:1202.3129](https://arxiv.org/abs/1202.3129)
- [10] Payne, S.: *Boundary complexes and weight filtrations*. [arXiv:1109.4286](https://arxiv.org/abs/1109.4286)
- [11] Saito, M.: *Hodge Modules*, Publ. Res. Inst. Math. Sci. 26 (1990), no. 2, 221-333.
- [12] Weber, A.: *Pure homology of algebraic varieties*. Topology Vol. 43, no 3, (2004) p. 635-644. DOI: [10.1016/j.top.2003.09.001](https://doi.org/10.1016/j.top.2003.09.001)
- [13] Weber, A.: *Purity at the end*. Preprint 2009, <http://arxiv.org/abs/0809.1885>

DEPARTMENT OF MATHEMATICS OF WARSAW UNIVERSITY, BANACHA 2, 02-097, WARSZAWA, POLAND  
E-mail address: [aweber@mimuw.edu.pl](mailto:aweber@mimuw.edu.pl)