
HOMOLOGICAL REPRESENTATIONS OF BRAID GROUPS AND KZ CONNECTIONS

TOSHITAKE KOHNO

ABSTRACT. We give a relation between the homological representations of the braid groups studied by Lawrence, Krammer and Bigelow and the monodromy of the KZ equation with values in the space of null vectors in the tensor product of Verma modules of $sl_2(\mathbf{C})$ when the parameters are generic. We also discuss the case of special parameters in relation with integral representations of the space of conformal blocks by hypergeometric integrals.

1. INTRODUCTION

The purpose of this article is to review recent developments concerning the relation between homological representations of the braid groups and the monodromy representations of the Knizhnik-Zamolodchikov (KZ) connections.

Homological representations appeared in the work of Lawrence [15] in relation with Hecke algebra representations of the braid groups. These representations were extensively investigated by Bigelow and Krammer and are called the Lawrence-Krammer-Bigelow (LKB) representations. In particular, they independently proved in [2] and [14] that the LKB representations of the braid groups are faithful.

On the other hand, it was shown by Schechtman-Varchenko [19] and others that the solutions of the KZ equation are expressed by hypergeometric integrals. By means of these integral representations we describe a relation between the LKB representations and the monodromy representations of the KZ connections. There are two parameters λ and κ , which are related to the highest weight and the KZ connection respectively. We consider the KZ equation with values in the space of null vectors in the tensor product of Verma modules of $sl_2(\mathbf{C})$ and show that a specialization of the LKB representation is equivalent to the monodromy representation of such KZ equation for generic parameters λ and κ . A more detailed treatment of this result is given in [13]. A relation between the Krammer representations of Artin groups and the monodromy of KZ connections is also described by an infinitesimal method by Marin [17].

The case of special parameters are important from the viewpoint of conformal field theory (see [6], [20] and [22]). We discuss the problem of expressing the space of conformal blocks by hypergeometric integrals. It is important to understand the KZ connection for the conformal block bundle as a Gauss-Manin connection. We refer the reader to the work of Looijenga [16] in recent advances concerning this subject.

The paper is organized in the following way. In Section 2 we recall the definition of the LKB representations. In Section 3 we describe the KZ connection. In Section 4 we review basic properties of Verma modules for the Lie algebra $sl_2(\mathbf{C})$ and define the space of null vectors. In Section 5 we state a comparison theorem between the LKB representations and the monodromy representations of the KZ connections with values in the space of null vectors. In Section 6 we describe the horizontal sections for the KZ connections by means of hypergeometric integrals. Section 7 is devoted to a proof of the comparison theorem stated in Section 5. We discuss the case of special parameters in relation with the space of conformal blocks in Section 8.

2. LAWRENCE-KRAMMER-BIGELOW REPRESENTATIONS

For a space X we denote by $\mathcal{F}_n(X)$ the configuration space of ordered distinct n points in X . Namely,

$$\mathcal{F}_n(X) = \{(x_1, \dots, x_n) \in X^n ; x_i \neq x_j \text{ if } i \neq j\},$$

where X^n denotes the n -fold Cartesian product of X .

We denote by B_n the braid group with n strands. We fix a set of distinct n points in \mathbf{R}^2 as

$$Q = \{(1, 0), \dots, (n, 0)\},$$

where we set $p_\ell = (\ell, 0)$, $\ell = 1, \dots, n$. We take a 2-dimensional disk in \mathbf{R}^2 containing Q in the interior. We fix a positive integer m and consider the configuration space $\mathcal{F}_m(\Sigma)$ of ordered distinct m points in $\Sigma = D \setminus Q$, which is also denoted by $\mathcal{F}_{n,m}(D)$. The symmetric group \mathfrak{S}_m acts freely on $\mathcal{F}_m(\Sigma)$ by the permutations of distinct m points. The quotient space of $\mathcal{F}_m(\Sigma)$ by this action is by definition the configuration space of unordered distinct m points in Σ and is denoted by $\mathcal{C}_m(\Sigma)$. We also denote this configuration space by $\mathcal{C}_{n,m}(D)$.

In the papers by Bigelow [2], [3] and by Krammer [14] the case $m = 2$ was extensively studied, but for our purpose it is convenient to consider the case when m is an arbitrary positive integer such that $m \geq 2$.

We identify \mathbf{R}^2 with the complex plane \mathbf{C} . The quotient space $\mathbf{C}^m/\mathfrak{S}_m$ defined by the action of \mathfrak{S}_m by the permutations of coordinates is analytically isomorphic to \mathbf{C}^m by means of the elementary symmetric polynomials. Now the image of the hyperplanes defined by $t_i = p_\ell$, $\ell = 1, \dots, n$, and the diagonal hyperplanes $t_i = t_j$, $1 \leq i \leq j \leq m$, are complex codimension one irreducible subvarieties of the quotient space D^m/\mathfrak{S}_m . This allows us to give a description of the first homology group of $\mathcal{C}_{n,m}(D)$ as

$$(2.1) \quad H_1(\mathcal{C}_{n,m}(D); \mathbf{Z}) \cong \mathbf{Z}^{\oplus n} \oplus \mathbf{Z}$$

where the first n components correspond to the normal loops of the images of hyperplanes $t_i = p_\ell$, $\ell = 1, \dots, n$, and the last component corresponds to the normal loop of the image of the diagonal hyperplanes $t_i = t_j$, $1 \leq i \leq j \leq m$, namely, the discriminant set. We consider the homomorphism

$$(2.2) \quad \alpha : H_1(\mathcal{C}_{n,m}(D); \mathbf{Z}) \longrightarrow \mathbf{Z} \oplus \mathbf{Z}$$

defined by $\alpha(x_1, \dots, x_n, y) = (x_1 + \dots + x_n, y)$. Composing with the abelianization map $\pi_1(\mathcal{C}_{n,m}(D), x_0) \rightarrow H_1(\mathcal{C}_{n,m}(D); \mathbf{Z})$, we obtain the homomorphism

$$(2.3) \quad \beta : \pi_1(\mathcal{C}_{n,m}(D), x_0) \longrightarrow \mathbf{Z} \oplus \mathbf{Z}.$$

Let $\pi : \tilde{\mathcal{C}}_{n,m}(D) \rightarrow \mathcal{C}_{n,m}(D)$ be the covering corresponding to $\text{Ker } \beta$. Now the group $\mathbf{Z} \oplus \mathbf{Z}$ acts as the deck transformations of the covering π and the homology group $H_*(\tilde{\mathcal{C}}_{n,m}(D); \mathbf{Z})$ is considered to be a $\mathbf{Z}[\mathbf{Z} \oplus \mathbf{Z}]$ -module. Here $\mathbf{Z}[\mathbf{Z} \oplus \mathbf{Z}]$ stands for the group ring of $\mathbf{Z} \oplus \mathbf{Z}$. We express $\mathbf{Z}[\mathbf{Z} \oplus \mathbf{Z}]$ as the ring of Laurent polynomials $R = \mathbf{Z}[q^{\pm 1}, t^{\pm 1}]$. We consider the homology group

$$H_{n,m} = H_m(\tilde{\mathcal{C}}_{n,m}(D); \mathbf{Z})$$

as an R -module by the action of the deck transformations.

In a similar way as is described in the case of $m = 2$ in [2], it can be shown that $H_{n,m}$ is a free R -module of rank

$$(2.4) \quad d_{n,m} = \binom{m+n-2}{m}.$$

A basis of $H_{n,m}$ as a free R -module is discussed in relation with the homology of local systems in the next sections. Let $\mathcal{M}(D, Q)$ denote the mapping class group of the pair (D, Q) ,

which consists of the isotopy classes of homeomorphisms of D which fix Q setwise and fix the boundary ∂D pointwise. The braid group B_n is naturally isomorphic to the mapping class group $\mathcal{M}(D, Q)$. Now a homeomorphism f representing a class in $\mathcal{M}(D, Q)$ induces a homeomorphism $\tilde{f} : \mathcal{C}_{n,m}(D) \rightarrow \mathcal{C}_{n,m}(D)$, which is uniquely lifted to a homeomorphism of $\tilde{\mathcal{C}}_{n,m}(D)$. This homeomorphism commutes with the deck transformations.

Therefore, for $m \geq 2$ we obtain a representation of the braid group

$$(2.5) \quad \rho_{n,m} : B_n \longrightarrow \text{Aut}_R H_{n,m}$$

which is called the homological representation of the braid group or the Lawrence-Krammer-Bigelow (LKB) representation. Let us remark that in the case $m = 1$ the above construction gives the reduced Burau representation over $\mathbf{Z}[q^{\pm 1}]$.

3. KZ CONNECTION

Let \mathfrak{g} be a complex semi-simple Lie algebra and $\{I_\mu\}$ be an orthonormal basis of \mathfrak{g} with respect to the Cartan-Killing form. Although we deal with the case $\mathfrak{g} = \mathfrak{sl}_2(\mathbf{C})$, we formulate the KZ connection in the case of a general complex semi-simple Lie algebra. We set $\Omega = \sum_\mu I_\mu \otimes I_\mu$. Let $r_i : \mathfrak{g} \rightarrow \text{End}(V_i)$, $1 \leq i \leq n$, be representations of the Lie algebra \mathfrak{g} . We denote by Ω_{ij} the action of Ω on the i -th and j -th components of the tensor product $V_1 \otimes \cdots \otimes V_n$. It is known that the Casimir element $c = \sum_\mu I_\mu \cdot I_\mu$ lies in the center of the universal enveloping algebra $U\mathfrak{g}$. Let us denote by $\Delta : U\mathfrak{g} \rightarrow U\mathfrak{g} \otimes U\mathfrak{g}$ the coproduct, which is defined to be the algebra homomorphism determined by $\Delta(x) = x \otimes 1 + 1 \otimes x$ for $x \in \mathfrak{g}$. Since Ω is expressed as $\Omega = \frac{1}{2}(\Delta(c) - c \otimes 1 - 1 \otimes c)$ we have the relation

$$(3.1) \quad [\Omega, x \otimes 1 + 1 \otimes x] = 0$$

for any $x \in \mathfrak{g}$ in the tensor product $U\mathfrak{g} \otimes U\mathfrak{g}$. By means of the above relation we obtain the quadratic relations:

$$(3.2) \quad [\Omega_{ik}, \Omega_{ij} + \Omega_{jk}] = 0, \quad (i, j, k \text{ distinct}),$$

$$(3.3) \quad [\Omega_{ij}, \Omega_{k\ell}], \quad (i, j, k, \ell \text{ distinct}),$$

which are called the infinitesimal pure braid relations.

We denote by X_n the configuration space $\mathcal{F}_n(\mathbf{C})$. Namely, X_n is defined as

$$X_n = \{(z_1, \dots, z_n) \in \mathbf{C}^n ; z_i \neq z_j \text{ if } i \neq j\}.$$

We define the Knizhnik-Zamolodchikov (KZ) connection as the 1-form

$$(3.4) \quad \omega = \frac{1}{\kappa} \sum_{1 \leq i < j \leq n} \Omega_{ij} d \log(z_i - z_j)$$

with values in $\text{End}(V_1 \otimes \cdots \otimes V_n)$ for a non-zero complex parameter κ . We have $d\omega = 0$ on X_n .

We set $\omega_{ij} = d \log(z_i - z_j)$, $1 \leq i \neq j \leq n$. It follows from the above infinitesimal pure braid relations among Ω_{ij} together with Arnold's relation

$$(3.5) \quad \omega_{ij} \wedge \omega_{jk} + \omega_{jk} \wedge \omega_{k\ell} + \omega_{k\ell} \wedge \omega_{ij} = 0$$

that $\omega \wedge \omega = 0$ holds. It follows that ω defines a flat connection for a trivial vector bundle over the configuration space X_n with fiber $V_1 \otimes \cdots \otimes V_n$. A horizontal section of the above flat bundle is a solution of the total differential equation

$$(3.6) \quad d\varphi = \omega\varphi$$

for a function $\varphi(z_1, \dots, z_n)$ on X_n with values in $V_1 \otimes \dots \otimes V_n$. This total differential equation can be expressed as a system of partial differential equations

$$(3.7) \quad \frac{\partial \varphi}{\partial z_i} = \frac{1}{\kappa} \sum_{j, j \neq i} \frac{\Omega_{ij}}{z_i - z_j} \varphi, \quad 1 \leq i \leq n,$$

which is called the KZ equation. The KZ equation was first introduced in [7] as the differential equation satisfied by n -point functions in Wess-Zumino-Witten conformal field theory on the Riemann sphere with the gauge symmetry of affine Lie algebras.

Let $\phi(z_1, \dots, z_n)$ be the matrix whose columns are linearly independent solutions of the KZ equation. By considering the analytic continuation of the solutions with respect to a loop γ in X_n with a base point x_0 we obtain the matrix $\theta(\gamma)$ defined by

$$(3.8) \quad \phi(z_1, \dots, z_n) \mapsto \phi(z_1, \dots, z_n) \theta(\gamma).$$

Since the KZ connection ω is flat the matrix $\theta(\gamma)$ depends only on the homotopy class of γ . The fundamental group $\pi_1(X_n, x_0)$ is the pure braid group P_n . As the above holonomy of the connection ω we have linear representations of the pure braid group

$$(3.9) \quad \theta : P_n \rightarrow \text{GL}(V_1 \otimes \dots \otimes V_n)$$

depending on the parameter κ .

The symmetric group \mathfrak{S}_n acts on X_n by the permutations of coordinates. We denote the quotient space X_n/\mathfrak{S}_n by Y_n . The fundamental group of Y_n is the braid group B_n . In the case $V_1 = \dots = V_n = V$, the symmetric group \mathfrak{S}_n acts diagonally on the trivial vector bundle over X_n with fiber $V^{\otimes n}$ and the connection ω is invariant by this action. Thus we have a one-parameter family of linear representations of the braid group

$$(3.10) \quad \theta : B_n \rightarrow \text{GL}(V^{\otimes n}).$$

It is known by [5] and [10] that this representation is described by means of quantum groups.

4. SPACE OF NULL VECTORS

Let us recall basic definitions about the Lie algebra $sl_2(\mathbf{C})$ and its Verma modules. As a complex vector space the Lie algebra $sl_2(\mathbf{C})$ has a basis H, E and F satisfying the relations:

$$(4.1) \quad [H, E] = 2E, \quad [H, F] = -2F, \quad [E, F] = H.$$

For a complex number λ we denote by M_λ the Verma module of $sl_2(\mathbf{C})$ with highest weight λ . Namely, there is a non-zero vector $v_\lambda \in M_\lambda$ called the highest weight vector satisfying

$$(4.2) \quad H v_\lambda = \lambda v_\lambda, \quad E v_\lambda = 0$$

and M_λ is spanned by $F^j v_\lambda, j \geq 0$. The elements H, E and F act on this basis as

$$(4.3) \quad \begin{cases} H \cdot F^j v_\lambda = (\lambda - 2j) F^j v_\lambda \\ E \cdot F^j v_\lambda = j(\lambda - j + 1) F^{j-1} v_\lambda \\ F \cdot F^j v_\lambda = F^{j+1} v_\lambda. \end{cases}$$

It is known that if $\lambda \in \mathbf{C}$ is not a non-negative integer, then the Verma module M_λ is irreducible.

The Shapovalov form is the symmetric bilinear form

$$S : M_\lambda \times M_\lambda \longrightarrow \mathbf{C}$$

characterized by the conditions:

$$\begin{aligned} S(v, v) &= 1, \\ S(Fx, y) &= S(x, Ey) \quad \text{for any } x, y \in M_\lambda. \end{aligned}$$

It follows from the above conditions that

$$\begin{aligned} S(F^i v, F^j v) &= 0 \text{ if } i \neq j, \\ S(F^i v, F^i v) &= i! \lambda(\lambda - 1) \cdots (\lambda - i + 1). \end{aligned}$$

For $\Lambda = (\lambda_1, \dots, \lambda_n) \in \mathbf{C}^n$ we put $|\Lambda| = \lambda_1 + \dots + \lambda_n$ and consider the tensor product $M_{\lambda_1} \otimes \dots \otimes M_{\lambda_n}$. For a non-negative integer m we define the space of weight vectors with weight $|\Lambda| - 2m$ by

$$(4.4) \quad W[|\Lambda| - 2m] = \{x \in M_{\lambda_1} \otimes \dots \otimes M_{\lambda_n} ; Hx = (|\Lambda| - 2m)x\}$$

and consider the space of null vectors defined by

$$(4.5) \quad N[|\Lambda| - 2m] = \{x \in W[|\Lambda| - 2m] ; Ex = 0\}.$$

The KZ connection ω defined in the previous section commutes with the diagonal action of \mathfrak{g} on $V_{\lambda_1} \otimes \dots \otimes V_{\lambda_n}$, hence it acts on the space of null vectors $N[|\Lambda| - 2m]$. This means that it is sensible to investigate the horizontal sections of the KZ connection with values in $N[|\Lambda| - 2m]$.

We also consider the space of coinvariant tensors defined by

$$L[|\Lambda| - 2m] = W[|\Lambda| - 2m] / (F \cdot W[|\Lambda| - 2m + 2]).$$

The dual space $L[|\Lambda| - 2m]^*$ is identified with the space of linear forms $f : W[|\Lambda| - 2m] \rightarrow \mathbf{C}$ such that $f(x) = 0$ for $x \in F \cdot W[|\Lambda| - 2m + 2]$. The isomorphism $M_\lambda \cong M_\lambda^*$ given by the Shapovalov form induces an isomorphism

$$N[|\Lambda| - 2m] \cong L[|\Lambda| - 2m]^*.$$

5. RELATION BETWEEN LKB REPRESENTATION AND KZ CONNECTION

We fix a complex number λ and consider the case $\lambda_1 = \dots = \lambda_n = \lambda$ and deal with the space of null vectors

$$N[n\lambda - 2m] \subset M_\lambda^{\otimes n}.$$

As the monodromy of the KZ connection

$$\omega = \frac{1}{\kappa} \sum_{1 \leq i < j \leq n} \Omega_{ij} d \log(z_i - z_j)$$

with values in $N[n\lambda - 2m]$ we obtain the linear representation of the braid group

$$\theta_{\lambda, \kappa} : B_n \rightarrow \text{Aut } N[n\lambda - 2m].$$

The next theorem describes a relationship between a specialization of the Lawrence-Krammer-Bigelow representation $\rho_{n,m}$ and the representation $\theta_{\lambda, \kappa}$.

Theorem 5.1. *There exists an open dense subset U in $(\mathbf{C}^*)^2$ such that for $(\lambda, \kappa) \in U$ the Lawrence-Krammer-Bigelow representation $\rho_{n,m}$ with the specialization*

$$q = e^{-2\pi\sqrt{-1}\lambda/\kappa}, \quad t = e^{2\pi\sqrt{-1}/\kappa}$$

is equivalent to the monodromy representation of the KZ connection $\theta_{\lambda, \kappa}$ with values in the space of null vectors

$$N[n\lambda - 2m] \subset M_\lambda^{\otimes n}.$$

In the rest of this section we recall some basic notions needed to prove Theorem 5.1. First, we recall some basic definition for local systems. Let M be a smooth manifold and V a complex vector space. Given a linear representation of the fundamental group

$$r : \pi_1(M, x_0) \longrightarrow GL(V)$$

there is an associated flat vector bundle E over M . The local system \mathcal{L} associated to the representation r is the sheaf of horizontal sections of the flat bundle E . Let $\pi : \widetilde{M} \rightarrow M$ be the universal covering. We denote by $\mathbf{Z}\pi_1$ the group ring of the fundamental group $\pi_1(M, x_0)$. We consider the chain complex

$$C_*(\widetilde{M}) \otimes_{\mathbf{Z}\pi_1} V$$

with the boundary map defined by $\partial(c \otimes v) = \partial c \otimes v$. Here $\mathbf{Z}\pi_1$ acts on $C_*(\widetilde{M})$ via the deck transformations and on V via the representation r . The homology of this chain complex is called the homology of M with coefficients in the local system \mathcal{L} and is denoted by $H_*(M, \mathcal{L})$.

The open dense set in Theorem 5.1 is closely related to the vanishing theorem of the homology of generic local system over the complement of hyperplane arrangements. Let us briefly recall this aspect. Let $\mathcal{A} = \{H_1, \dots, H_N\}$ be a set of affine hyperplanes in the complex vector space \mathbf{C}^n . We call the set \mathcal{A} a complex hyperplane arrangement. We consider the complement

$$M(\mathcal{A}) = \mathbf{C}^n \setminus \bigcup_{H \in \mathcal{A}} H.$$

Let \mathcal{L} be a complex rank one local system over $M(\mathcal{A})$ associated with a representation of the fundamental group

$$r : \pi_1(M(\mathcal{A}), x_0) \longrightarrow \mathbf{C}^*.$$

We investigate the homology of $M(\mathcal{A})$ with coefficients in the local system \mathcal{L} . For our purpose the homology of locally finite chains $H_*^{lf}(M(\mathcal{A}), \mathcal{L})$ also plays an important role.

We summarize basic properties of the above homology groups. For a complex hyperplane arrangement \mathcal{A} we choose a smooth compactification $i : M(\mathcal{A}) \rightarrow X$ with normal crossing divisors. We shall say that the local system \mathcal{L} is generic if and only if an isomorphism

$$(5.1) \quad i_*\mathcal{L} \cong i_!\mathcal{L}$$

holds, where $i_*\mathcal{L}$ is the direct image and $i_!\mathcal{L}$ is the extension of \mathcal{L} by 0 outside of $M(\mathcal{A})$ as a sheaf. The condition $i_*\mathcal{L} \cong i_!\mathcal{L}$ means that the monodromy of \mathcal{L} along any divisor at infinity is not equal to 1. The following theorem was shown in [9].

Theorem 5.2. *If the local system \mathcal{L} is generic in the above sense, then there is an isomorphism*

$$H_*(M(\mathcal{A}), \mathcal{L}) \cong H_*^{lf}(M(\mathcal{A}), \mathcal{L}).$$

Moreover, we have $H_k(M(\mathcal{A}), \mathcal{L}) = 0$ for any $k \neq n$.

Proof. In general we have isomorphisms

$$H^*(X, i_*\mathcal{L}) \cong H^*(M(\mathcal{A}), \mathcal{L}), \quad H^*(X, i_!\mathcal{L}) \cong H_c^*(M(\mathcal{A}), \mathcal{L})$$

where H_c denotes cohomology with compact supports.

There are Poincaré duality isomorphisms:

$$\begin{aligned} H_k^{lf}(M(\mathcal{A}), \mathcal{L}) &\cong H^{2n-k}(M(\mathcal{A}), \mathcal{L}), \\ H_k(M(\mathcal{A}), \mathcal{L}) &\cong H_c^{2n-k}(M(\mathcal{A}), \mathcal{L}). \end{aligned}$$

By the hypothesis $i_*\mathcal{L} \cong i_!\mathcal{L}$ we obtain an isomorphism

$$H_k^{lf}(M(\mathcal{A}), \mathcal{L}) \cong H_k(M(\mathcal{A}), \mathcal{L}).$$

It follows from the above Poincaré duality isomorphisms and the fact that $M(\mathcal{A})$ has a homotopy type of a CW complex of dimension at most n we have

$$\begin{aligned} H_k^{lf}(M(\mathcal{A}), \mathcal{L}) &\cong 0, & k < n \\ H_k(M(\mathcal{A}), \mathcal{L}) &\cong 0, & k > n. \end{aligned}$$

Therefore we obtain $H_k(M(\mathcal{A}), \mathcal{L}) = 0$ for any $k \neq n$. \square

For a positive integer m we consider the projection map

$$(5.2) \quad \pi_{n,m} : X_{n+m} \longrightarrow X_n$$

given by $\pi_{n,m}(z_1, \dots, z_n, t_1, \dots, t_m) = (z_1, \dots, z_n)$, which defines a fiber bundle over X_n . For $p \in X_n$ the fiber $\pi_{n,m}^{-1}(p)$ is denoted by $X_{n,m}$. Let (p_1, \dots, p_n) be the coordinates for p . Then, $X_{n,m}$ is the complement of hyperplanes defined by

$$(5.3) \quad t_i = p_\ell, \quad 1 \leq i \leq m, \quad 1 \leq \ell \leq n, \quad t_i = t_j, \quad 1 \leq i < j \leq m.$$

Such arrangement of hyperplanes is called a discriminantal arrangement. The symmetric group \mathfrak{S}_m acts on $X_{n,m}$ by the permutations of the coordinates functions t_1, \dots, t_m . We put $Y_{n,m} = X_{n,m}/\mathfrak{S}_m$.

Identifying \mathbf{R}^2 with the complex plane \mathbf{C} , we have the inclusion map

$$(5.4) \quad \iota : \mathcal{F}_{n,m}(D) \longrightarrow X_{n,m},$$

which is a homotopy equivalence. By taking the quotient by the action of the symmetric group \mathfrak{S}_m , we have the inclusion map

$$(5.5) \quad \bar{\iota} : \mathcal{C}_{n,m}(D) \longrightarrow Y_{n,m},$$

which is also a homotopy equivalence.

We take $p = (1, 2, \dots, n)$ as a base point. We consider a local system over $X_{n,m}$ defined in the following way. Let $\xi_{i\ell}$ and η_{ij} be normal loops around the hyperplanes $t_i = p_\ell$ and $t_i = t_j$ respectively. We fix complex numbers α_ℓ , $1 \leq \ell \leq n$, and γ and by the correspondence

$$\xi_{i\ell} \mapsto e^{2\pi\sqrt{-1}\alpha_\ell}, \quad \eta_{ij} \mapsto e^{4\pi\sqrt{-1}\gamma}$$

we obtain the representation

$$r : \pi_1(X_{n,m}, x_0) \longrightarrow \mathbf{C}^*.$$

We denote by \mathcal{L} the associated rank one local system on $X_{n,m}$.

Let us consider the embedding

$$(5.6) \quad i_0 : X_{n,m} \longrightarrow (\mathbf{C}P^1)^m = \underbrace{\mathbf{C}P^1 \times \dots \times \mathbf{C}P^1}_m.$$

Then we take blowing-ups at multiple points $\pi : (\widehat{\mathbf{C}P^1})^m \longrightarrow (\mathbf{C}P^1)^m$ and obtain a smooth compactification $i : X_{n,m} \rightarrow (\widehat{\mathbf{C}P^1})^m$ with normal crossing divisors. We can write down the condition $i_*\mathcal{L} \cong i_!\mathcal{L}$ explicitly by computing the monodromy of the local system \mathcal{L} along divisors at infinity.

The local system \mathcal{L} on $X_{n,m}$ is invariant under the action of the symmetric group \mathfrak{S}_m and induces the local system $\bar{\mathcal{L}}$ on $Y_{n,m}$. We will deal with the case $\alpha_1 = \dots = \alpha_\ell = \alpha$. By a similar argument using a smooth compactification of $Y_{n,m}$ we have the following proposition.

Proposition 5.1. *There is an open dense subset V in \mathbf{C}^2 such that for $(\alpha, \gamma) \in V$ the associated local system $\bar{\mathcal{L}}$ on $Y_{n,m}$ satisfies*

$$H_*(Y_{n,m}, \bar{\mathcal{L}}) \cong H_*^{lf}(Y_{n,m}, \bar{\mathcal{L}})$$

and $H_k(Y_{n,m}, \overline{\mathcal{L}}) = 0$ for any $k \neq m$. Moreover, we have

$$(5.7) \quad \dim H_m(Y_{n,m}, \overline{\mathcal{L}}^*) = d_{n,m},$$

where we use the same notation as in equation (2.4) for $d_{n,m}$.

For the purpose of describing the homology groups $H_m^{lf}(X_{n,m}, \mathcal{L})$ and $H_m^{lf}(Y_{n,m}, \overline{\mathcal{L}})$ we introduce the following notation. We fix the base point $p = (1, \dots, n)$. For non-negative integers m_1, \dots, m_{n-1} satisfying

$$(5.8) \quad m_1 + \dots + m_{n-1} = m$$

we define a bounded chamber $\Delta_{m_1, \dots, m_{n-1}}$ in \mathbf{R}^m by

$$\begin{aligned} 1 &< t_1 < \dots < t_{m_1} < 2 \\ 2 &< t_{m_1+1} < \dots < t_{m_1+m_2} < 3 \\ &\dots \\ n-1 &< t_{m_1+\dots+m_{n-2}+1} + \dots + t_m < n. \end{aligned}$$

We put $M = (m_1, \dots, m_{n-1})$ and we write Δ_M for $\Delta_{m_1, \dots, m_{n-1}}$. We denote by $\overline{\Delta}_M$ the image of Δ_M by the projection map $\pi_{n,m}$. The bounded chamber Δ_M defines a homology class $[\Delta_M] \in H_m^{lf}(X_{n,m}, \mathcal{L})$ and its image $\overline{\Delta}_M$ defines a homology class $[\overline{\Delta}_M] \in H_m^{lf}(Y_{n,m}, \overline{\mathcal{L}})$. We shall observe that under certain generic conditions $[\overline{\Delta}_M]$ for $M = (m_1, \dots, m_{n-1})$ with $m_1 + \dots + m_{n-1} = m$ form a basis of $H_m^{lf}(Y_{n,m}, \overline{\mathcal{L}})$.

As we have shown in Theorem 5.2 there is an isomorphism $H_m(X_{n,m}, \mathcal{L}) \cong H_m^{lf}(X_{n,m}, \mathcal{L})$ if the condition $i_*\mathcal{L} \cong i_!\mathcal{L}$ is satisfied. In this situation we denote by $[\tilde{\Delta}_M]$ the homology class in $H_m(X_{n,m}, \mathcal{L})$ corresponding to $[\Delta_M]$ in the above isomorphism and call $[\tilde{\Delta}_M]$ the regularized cycle for $[\Delta_M]$.

6. HYPERGEOMETRIC INTEGRALS

In this section we describe solutions of the KZ equation for the case $\mathfrak{g} = \mathfrak{sl}_2(\mathbf{C})$ by means of hypergeometric integrals following Schechtman and Varchenko [19]. A description of the solutions of the KZ equation was also given by Date, Jimbo, Matsuo and Miwa [4]. We refer the reader to [1] and [18] for general treatments of hypergeometric integrals.

For parameters κ and λ we consider the multi-valued function

$$(6.1) \quad \Phi_{n,m} = \prod_{1 \leq i < j \leq n} (z_i - z_j)^{\frac{\lambda_i \lambda_j}{\kappa}} \prod_{1 \leq i \leq m, 1 \leq \ell \leq n} (t_i - z_\ell)^{-\frac{\lambda_\ell}{\kappa}} \prod_{1 \leq i < j \leq m} (t_i - t_j)^{\frac{2}{\kappa}}$$

defined over X_{n+m} . Let \mathcal{L} denote the local system associated to the multi-valued function Φ . The restriction of \mathcal{L} on the fiber $X_{n,m}$ is the local system associated with the parameters

$$(6.2) \quad \alpha_\ell = -\frac{\lambda_\ell}{\kappa}, \quad 1 \leq \ell \leq n, \quad \gamma = \frac{1}{\kappa}.$$

The symmetric group \mathfrak{S}_m acts on $X_{n,m}$ by the permutations of the coordinate functions t_1, \dots, t_m . The function $\Phi_{n,m}$ is invariant by the action of \mathfrak{S}_m . The local system \mathcal{L} over $X_{n,m}$ defines a local system on $Y_{n,m}$, which we denote by $\overline{\mathcal{L}}$. The local system dual to \mathcal{L} is denoted by \mathcal{L}^* .

We put $v = v_{\lambda_1} \otimes \dots \otimes v_{\lambda_n}$ and for $J = (j_1, \dots, j_n)$ set $F^J v = F^{j_1} v_{\lambda_1} \otimes \dots \otimes F^{j_n} v_{\lambda_n}$, where j_1, \dots, j_n are non-negative integers. The weight space $W[|\Lambda| - 2m]$ has a basis $F^J v$ for each J

with $|J| = j_1 + \cdots + j_n = m$. For the sequence of integers $(i_1, \dots, i_m) = (\underbrace{1, \dots, 1}_{j_1}, \dots, \underbrace{n, \dots, n}_{j_n})$

we set

$$(6.3) \quad S_J(z, t) = \frac{1}{(t_1 - z_{i_1}) \cdots (t_m - z_{i_m})}$$

and define the rational function $R_J(z, t)$ by

$$(6.4) \quad R_J(z, t) = \frac{1}{j_1! \cdots j_n!} \sum_{\sigma \in \mathfrak{S}_m} S_J(z_1, \dots, z_n, t_{\sigma(1)}, \dots, t_{\sigma(m)}).$$

For example, we have

$$\begin{aligned} R_{(1,0,\dots,0)}(z, t) &= \frac{1}{t_1 - z_1}, & R_{(2,0,\dots,0)}(z, t) &= \frac{1}{(t_1 - z_1)(t_2 - z_1)} \\ R_{(1,1,0,\dots,0)}(z, t) &= \frac{1}{(t_1 - z_1)(t_2 - z_2)} + \frac{1}{(t_2 - z_1)(t_1 - z_2)} \end{aligned}$$

and so on.

Since $\pi_{n,m} : X_{m+n} \rightarrow X_m$ is a fiber bundle with fiber $X_{n,m}$ the fundamental group of the base space X_n acts naturally on the homology group $H_m(X_{n,m}, \mathcal{L}^*)$. Thus we obtain a representation of the pure braid group

$$(6.5) \quad r_{n,m} : P_n \longrightarrow \text{Aut } H_m(X_{n,m}, \mathcal{L}^*)$$

which defines a local system on X_n denoted by $\mathcal{H}_{n,m}$. In the case $\lambda_1 = \cdots = \lambda_n$ there is a representation of the braid group

$$(6.6) \quad r_{n,m} : B_n \longrightarrow \text{Aut } H_m(Y_{n,m}, \overline{\mathcal{L}}^*)$$

which defines a local system $\overline{\mathcal{H}}_{n,m}$ on $Y_{n,m}$. For any horizontal section $c(z)$ of the local system $\mathcal{H}_{n,m}$ we consider the hypergeometric type integral

$$(6.7) \quad \int_{c(z)} \Phi_{n,m} R_J(z, t) dt_1 \wedge \cdots \wedge dt_m$$

for the above rational function $R_J(z, t)$.

According to Schechtman and Varchenko, solutions of the KZ equation are described in the following way.

Theorem 6.1 (Schechtman and Varchenko [19]). *The integral*

$$\sum_{|J|=m} \left(\int_{c(z)} \Phi_{n,m} R_J(z, t) dt_1 \wedge \cdots \wedge dt_m \right) F^J v$$

lies in the space of null vectors $N[|\Lambda| - 2m]$ and is a solution of the KZ equation.

We assume the conditions $i_* \mathcal{L} \cong i_1 \mathcal{L}$ and $i_* \overline{\mathcal{L}} \cong i_1 \overline{\mathcal{L}}$ in the following. These conditions are satisfied for (λ, κ) in an open dense subset in $(\mathbf{C}^*)^2$. By the assumption we have an isomorphism $H_m(X_{n,m}, \mathcal{L}) \cong H_m^{lf}(X_{n,m}, \mathcal{L})$ and we can take the regularized cycles $[\tilde{\Delta}_M] \in H_m(X_{n,m}, \mathcal{L})$ for the bounded chamber Δ_M .

We will consider the integral

$$\sum_{|J|=m} \left(\int_{\Delta_M} \Phi_{n,m} R_J(z, t) dt_1 \wedge \cdots \wedge dt_m \right) F^J v$$

in the space of null vectors $N[|\Lambda| - 2m]$. In general the above integral is divergent. We replace the integration cycle by the regularized cycle $[\tilde{\Delta}_M]$ to obtain the convergent integral. This is called the regularized integral. We refer the reader to [1] for details on this aspect.

Let us denote by $\Omega^p(X_{n,m})$ the space of smooth p -forms on $X_{n,m}$. The twisted de Rham complex $(\Omega^*(X_{n,m}), \nabla)$ is a complex with the differential

$$\nabla : \Omega^p(X_{n,m}) \longrightarrow \Omega^{p+1}(X_{n,m})$$

defined by

$$\nabla\varphi = d\varphi + d\log\Phi \wedge \varphi.$$

There is a non-degenerate pairing between the homology with local coefficients and the cohomology of the twisted de Rham complex

$$H_m(X_{n,m}, \mathcal{L}^*) \times H^m(\Omega^*(X_{n,m}), \nabla) \longrightarrow \mathbf{C}$$

given by

$$(c, \varphi) \mapsto \int_c \Phi\varphi.$$

We define the map $\rho : W[|\lambda| - 2m] \longrightarrow \Omega^*(X_{n,m})$ by

$$\rho(F^J v) = R_J(z, t) dt_1 \wedge \cdots \wedge dt_m$$

for J with $|J| = m$. It is shown in [6] such that ρ induces a map

$$L[|\lambda| - 2m] \longrightarrow H^m(\Omega^*(X_{n,m}), \nabla).$$

This means that $\rho(w)$ is a closed form for any $w \in W[|\lambda| - 2m]$ and that the image of $F \cdot W[|\lambda| - 2m + 2]$ is contained in the space of exact forms in the twisted de Rham complex $(\Omega^*(X_{n,m}), \nabla)$.

7. PROOF OF THEOREM 5.1

In this section we give a proof of Theorem 5.1. We have the following proposition.

Proposition 7.1. *There exists an open dense subset U in $(\mathbf{C}^*)^2$ such that for $(\lambda, \kappa) \in U$ the following properties (1) and (2) are satisfied.*

- (1) *The integrals in Theorem 6.1 over $[\tilde{\Delta}_M]$ for $M = (m_1, \dots, m_{n-1})$ with $m_1 + \dots + m_{n-1} = m$ are linearly independent.*
- (2) *The homology classes $[\tilde{\Delta}_M]$ for $M = (m_1, \dots, m_{n-1})$ with $m_1 + \dots + m_{n-1} = m$ form a basis of $H_m^{lf}(Y_{n,m}, \overline{\mathcal{L}}^*) \cong H_m(Y_{n,m}, \overline{\mathcal{L}}^*)$.*

Here m_1, \dots, m_{n-1} are non-negative integers.

Proof. We suppose that λ_1 is not a non-negative integer. Let us observe that for $\Lambda = (\lambda_1, \dots, \lambda_n)$ the space of null vectors $N[|\Lambda| - 2m]$ has dimension $d_{n,m}$. This can be shown as follows. First, let us consider the weight space

$$\begin{aligned} & (M_{\lambda_2} \otimes \cdots \otimes M_{\lambda_n})[\lambda_2 + \cdots + \lambda_n - 2m] \\ & = \{x \in M_{\lambda_2} \otimes \cdots \otimes M_{\lambda_n} ; Hx = (\lambda_2 + \cdots + \lambda_n - 2m)x\}. \end{aligned}$$

There is an isomorphism

$$\xi : (M_{\lambda_2} \otimes \cdots \otimes M_{\lambda_n})[\lambda_2 + \cdots + \lambda_n - 2m] \longrightarrow N[|\Lambda| - 2m]$$

defined by

$$u \mapsto v_{\lambda_1} \otimes u - \frac{1}{\lambda_1} Fv_{\lambda_1} \otimes Eu + \frac{1}{\lambda_1(\lambda_1 - 1)} F^2v_{\lambda_1} \otimes E^2u - \cdots$$

This shows that $N[|\Lambda| - 2m]$ has a basis indexed by $J' = (j_1, j_2, \dots, j_n)$ with $j_1 = 0$ and $j_2 + \cdots + j_n = m$, where j_2, \dots, j_n are non-negative integers. Let us denote by $S_{n,m}$ the set of

such indices J' . The above weight space has a basis $u_{J'}$ indexed by $J' \in S_{n,m}$. We have the corresponding basis $\xi(u_{J'})$ of $N[|\Lambda| - 2m]$.

We set $\alpha_1, \dots, \alpha_n$ and γ as in (6.2). We put

$$(7.1) \quad \tilde{\Phi}_{n,m} = \prod_{1 \leq i \leq m, 1 \leq \ell \leq n} (t_i - z_\ell)^{\alpha_\ell} \prod_{1 \leq i < j \leq m} (t_i - t_j)^{2\gamma}$$

and for $J' \in S_{n,m}$ put

$$(7.2) \quad \alpha'_{J'} = \prod_{k=2}^n (j_k)! \alpha_k (\alpha_k + \gamma) \cdots (\alpha_k + (j_k - 1)\gamma).$$

We assume that $\alpha_1, \dots, \alpha_n$ and γ are positive. We express the integral in Theorem 5.1 over the cycle Δ_M in the linear combination for the basis $\xi(u_{J'})$ of $N[|\Lambda| - 2m]$ and we denote by $\tilde{R}_{J'}(z, t)$ the corresponding rational function. In [21] Varchenko gave a formula for the determinant

$$(7.3) \quad \det_{M, J'} \left(\alpha_{J'} \int_{\Delta_M} \tilde{\Phi}_{n,m} \tilde{R}_{J'}(z, t) dt_1 \wedge \cdots \wedge dt_m \right),$$

where $M = (m_1, \dots, m_{n-1})$ with $m_1 + \cdots + m_{n-1} = m$ and $J' \in S_{n,m}$. According to Varchenko's formula the above determinant is expressed as a non-zero constant times the gamma factor given by

$$(7.4) \quad \prod_{i=0}^{m-1} \left(\frac{\Gamma((i+1)\gamma + 1)^{n-1}}{\Gamma(\gamma + 1)^{n-1}} \frac{\Gamma(\alpha_1 + i\gamma + 1) \cdots \Gamma(\alpha_n + i\gamma + 1)}{\Gamma(\alpha_1 + \cdots + \alpha_n + (2m - 2 - i)\gamma + 1)} \right)^{\nu_i}$$

where ν_i is defined by

$$(7.5) \quad \nu_i = \binom{m+n-i-3}{m-i-1}.$$

Since the gamma function does not have zeros and has only poles of order one at non-positive integers, it is clear that the determinant is zero only when the denominator of the gamma factor has a pole. Considering the regularized integrals over the cycles $[\tilde{\Delta}_M]$ we can analytically continue the determinant formula to complex numbers $\alpha_1, \dots, \alpha_n$ and γ .

Let us recall that we deal with the case

$$\alpha_\ell = -\frac{\lambda}{\kappa}, \quad 1 \leq \ell \leq n, \quad \gamma = \frac{1}{\kappa}.$$

From the determinant formula we observe that the linear independence for the solutions of the KZ equation in (1) in the statement of the proposition is satisfied for (λ, κ) in an open dense subset in $(\mathbf{C}^*)^2$. Under the same condition we have the linear independence for the homology classes $[\tilde{\Delta}_M]$ for $M = (m_1, \dots, m_{n-1})$ with $m_1 + \cdots + m_{n-1} = m$. Since we have $\dim H_m^{lf}(Y_{n,m}, \overline{\mathcal{L}}^*) = d_{m,n}$ we obtain the property (2). This completes the proof of our proposition. \square

Let us consider the specialization map

$$(7.6) \quad s : R = \mathbf{Z}[q^{\pm 1}, t^{\pm 1}] \longrightarrow \mathbf{C}$$

defined by the substitutions $q \mapsto e^{-2\pi\sqrt{-1}\lambda/\kappa}$ and $t \mapsto e^{2\pi\sqrt{-1}/\kappa}$. This induces in a natural way a homomorphism

$$(7.7) \quad H_m(\tilde{\mathcal{C}}_{n,m}(D); \mathbf{Z}) \longrightarrow H_m(Y_{n,m}, \overline{\mathcal{L}}^*).$$

We take a basis $[c_M]$ of $H_m(\tilde{\mathcal{C}}_{n,m}(D); \mathbf{Z})$ as the R -module for $M = (m_1, \dots, m_{n-1})$ with $m_1 + \cdots + m_{n-1} = m$ in such a way that $[c_M]$ maps to the regularized cycle for $[\tilde{\Delta}_M]$ by the above

specialization map. We observe that the LKB representation specialized at $q \mapsto e^{-2\pi\sqrt{-1}\lambda/\kappa}$ and $t \mapsto e^{2\pi\sqrt{-1}/\kappa}$ is identified with the linear representation of the braid group $r_{n,m} : B_n \rightarrow \text{Aut } H_m(Y_{n,m}, \overline{\mathcal{L}}^*)$.

Now the fundamental solutions of the KZ equation with values in $N[n\lambda - 2m]$ is given by the matrix of the form

$$\left(\int_{\tilde{\Delta}_M} \omega_{M'} \right)_{M, M'}$$

with $M = (m_1, \dots, m_{n-1})$ and $M' = (m'_1, \dots, m'_{n-1})$ such that $m_1 + \dots + m_{n-1} = m$ and $m'_1 + \dots + m'_{n-1} = m$. Here $\omega_{M'}$ is a multivalued m -form on $X_{n,m}$. The determinant of this matrix is computed in [22]. We observe that this determinant is non-zero for a generic (λ, κ) . This fact is used to show Proposition 7.1. The column vectors of the above matrix form a basis of the solutions of the KZ equation with values in $N[n\lambda - 2m]$. Thus the representation $r_{n,m} : B_n \rightarrow \text{Aut } H_m(Y_{n,m}, \overline{\mathcal{L}}^*)$ is equivalent to the action of B_n on the solutions of the KZ equation with values in $N[n\lambda - 2m]$. This completes the proof of Theorem 5.1.

As a consequence of the above argument we have an isomorphism

$$H_m(Y_{n,m}, \overline{\mathcal{L}}^*) \cong N[n\lambda - 2m],$$

which is equivariant under the action of the braid group B_n .

8. INTEGRAL REPRESENTATION OF THE SPACE OF CONFORMAL BLOCKS

Let us recall some basic definitions concerning the affine Lie algebras and their representations. We deal with the case $\mathfrak{g} = \mathfrak{sl}_2(\mathbf{C})$. Let us denote by $\mathbf{C}((\xi))$ the ring of formal Laurent series consisting of the power series $\sum_{n=-m}^{\infty} a_n \xi^n$, $a_n \in \mathbf{C}$ for some positive integer m . The loop algebra $L\mathfrak{g}$ is the tensor product $\mathfrak{g} \otimes \mathbf{C}((\xi))$ equipped with a Lie algebra structure by

$$[X \otimes f, Y \otimes g] = [X, Y] \otimes fg.$$

Now the affine Lie algebra $\widehat{\mathfrak{g}}$ is the central extension

$$\widehat{\mathfrak{g}} = L\mathfrak{g} \oplus \mathbf{C}c$$

with the Lie bracket defined by

$$[X \otimes f, Y \otimes g] = [X, Y] \otimes fg + \text{Res}_{\xi=0}(df \cdot g) \langle X, Y \rangle c$$

where $\langle X, Y \rangle$ stands for the Cartan-Killing form for \mathfrak{g} .

Let A_+ denote the subring of $\mathbf{C}((\xi))$ consisting of the power series of the form $\sum_{n>0} a_n \xi^n$. In a similar way, we denote by A_- the subring consisting of $\sum_{n<0} a_n \xi^n$. We put

$$\begin{aligned} N_+ &= [\mathfrak{g} \otimes A_+] \oplus \mathbf{C}E \\ N_0 &= \mathbf{C}H \oplus \mathbf{C}c \\ N_- &= [\mathfrak{g} \otimes A_-] \oplus \mathbf{C}F, \end{aligned}$$

which gives a direct sum decomposition of Lie algebras

$$\widehat{\mathfrak{g}} = N_+ \oplus N_0 \oplus N_-.$$

Let k and λ be complex numbers. We consider the representation $M_{k,\lambda}$ of $\widehat{\mathfrak{g}}$ with the non-zero vector v satisfying the following properties :

- (1) $N_+v = 0$, $Hv = \lambda v$, $cv = kv$
- (2) $M_{k,\lambda}$ is freely generated by v over $U(N_-)$,

where $U(N_-)$ is the universal enveloping algebra of N_- . The representation $M_{k,\lambda}$ is called the Verma module with highest weight λ and level k . It is known that for generic values k and λ the Verma module $M_{k,\lambda}$ is irreducible.

Now let us consider the case when k is a positive integer and λ is an integer such that $0 \leq \lambda \leq k$. In this case the Verma module $M_{k,\lambda}$ is not irreducible. In fact the vector $\chi = (E \otimes \xi)^{k-\lambda+1}v$ satisfies $N_+\chi = 0$ and $U(N_-)\chi$ is a proper submodule of $M_{k,\lambda}$. The quotient module $H_{k,\lambda}$ is irreducible and is called the integrable highest weight module with highest weight λ and level k . When we fix the level k we will write H_λ for $H_{k,\lambda}$. We refer the reader to Kac [8] for details.

We denote by V_λ the irreducible \mathfrak{g} -module with highest weight λ . Namely, there exists a non-zero vector $v \in V_\lambda$ such that $Hv = \lambda v$ and $Ev = 0$. The vector space V_λ has a basis $v, Fv, \dots, F^\lambda v$. The Verma module $M_{k,\lambda}$ is written as $U(N_-)V_\lambda$.

We define the space of conformal blocks for a Riemann sphere \mathbf{CP}^1 with marked points as follows. See [11] for a more detailed exposition of the subject. We take distinct points p_1, \dots, p_n, p_{n+1} such that $p_{n+1} = \infty$. We fix an affine coordinate function z for $\mathbf{CP}^1 \setminus \{\infty\}$. Let z_j be the coordinate for p_j , $1 \leq j \leq n$, and we set $\xi_j = z - z_j$. We take $\xi_{n+1} = 1/z$ as a local coordinate around ∞ . We assign integers $\lambda_1, \dots, \lambda_n, \lambda_{n+1}$ satisfying $0 \leq \lambda_j \leq k$, $1 \leq j \leq n+1$ to the points p_1, \dots, p_n, p_{n+1} . We set $p = (p_1, \dots, p_n, p_{n+1})$ and $\lambda = (\lambda_1, \dots, \lambda_n, \lambda_{n+1})$. Let us denote by \mathcal{M}_p the set of meromorphic functions on \mathbf{CP}^1 with poles at most at p_1, \dots, p_n, p_{n+1} .

The space of conformal blocks $\mathcal{H}(p, \lambda)$ is by definition the space of coinvariant tensors

$$(H_{\lambda_1} \otimes \dots \otimes H_{\lambda_{n+1}})/(\mathfrak{g} \otimes \mathcal{M}_p)$$

where $\mathfrak{g} \otimes \mathcal{M}_p$ acts diagonally on the tensor product $H_{\lambda_1} \otimes \dots \otimes H_{\lambda_{n+1}}$ by means of the Laurent expansion of a meromorphic function at p_1, \dots, p_{n+1} with respect to the local coordinates ξ_1, \dots, ξ_{n+1} . We define its dual space $\mathcal{H}(p, \lambda)^*$ as the space of multilinear forms

$$H_{\lambda_1} \otimes \dots \otimes H_{\lambda_{n+1}} \longrightarrow \mathbf{C}$$

invariant under the diagonal action of $\mathfrak{g} \otimes \mathcal{M}_p$ defined in the above way.

It turns out that the space of conformal blocks $\mathcal{H}(p, \lambda)$ is a finite dimensional complex vector space. In fact $\mathcal{H}(p, \lambda)$ is isomorphic to the quotient space of the space of coinvariant tensors defined by

$$(V_{\lambda_1} \otimes \dots \otimes V_{\lambda_{n+1}})/\mathfrak{g}$$

for the diagonal action of \mathfrak{g} . The kernel of the surjective homomorphism $(V_{\lambda_1} \otimes \dots \otimes V_{\lambda_{n+1}})/\mathfrak{g} \rightarrow \mathcal{H}(p, \lambda)$ is described by algebraic relations depending on z coming from the existence of the null vector χ in the definition of the integrable highest weight module H_λ . The dual construction gives an injective map

$$\mathcal{H}(p, \lambda)^* \longrightarrow \text{Hom}_{\mathfrak{g}}(V_{\lambda_1} \otimes \dots \otimes V_{\lambda_{n+1}}, \mathbf{C}).$$

We fix λ and consider the disjoint union

$$\mathcal{E}_\lambda = \bigcup_{(p_1, \dots, p_n) \in X_n} \mathcal{H}(p, \lambda),$$

which has a structure of a vector bundle over X_n . It turns out that the KZ connection with the parameter $\kappa = K + 2$ induces a flat connection on the vector bundle \mathcal{E}_λ . We call \mathcal{E}_λ the conformal block bundle. In a similar way, we define the dual conformal block bundle \mathcal{E}_λ^* over X_n whose fiber is $\mathcal{H}(p, \lambda)^*$.

In our case the space of coinvariant tensors $L[|\lambda| - 2m]$ with $|\lambda| = \lambda_1 + \dots + \lambda_n$ and $m = \frac{1}{2}(|\lambda| - \lambda_{n+1})$ is isomorphic to $(V_{\lambda_1} \otimes \dots \otimes V_{\lambda_{n+1}})/\mathfrak{g}$. We have the following theorem.

Theorem 8.1 (Feigin, Schechtman and Varchenko [6]). *The map*

$$\rho : L[|\lambda| - 2m] \rightarrow H^m(\Omega^m(X_{n,m}), \nabla)$$

factors through the projection map $L[|\lambda| - 2m] \rightarrow \mathcal{H}(p, \lambda)$.

Furthermore, as is stated in [6], it is shown in [22] that the induced map

$$\bar{\rho} : \mathcal{H}(p, \lambda) \longrightarrow H^m(\Omega^m(X_{n,m}), \nabla)$$

is injective. Let us consider the dual map $\bar{\phi} : H_m(X_{n,m}, \mathcal{L}^*) \longrightarrow \mathcal{H}(p, \lambda)^*$ defined by

$$\langle \bar{\phi}(c), w \rangle = \int_c \Phi \bar{\rho}(w)$$

for $w \in V_{\lambda_1} \otimes \cdots \otimes V_{\lambda_{n+1}}$. It follows from the above construction that the map $\bar{\phi}$ is surjective.

There is a flat bundle $\mathcal{H}_{n,m}$ over the configuration space X_n whose fiber is the homology $H_m(X_{n,m}, \mathcal{L}^*)$. We have a surjective bundle map

$$\mathcal{H}_{n,m} \longrightarrow \mathcal{E}_\lambda^*$$

which is compatible with the flat structures. It turns out that any horizontal section of the dual conformal block bundle \mathcal{E}_λ^* is expressed as

$$\langle \bar{\phi}(c(z)), w \rangle = \int_{c(z)} \Phi \bar{\rho}(w)$$

with a horizontal section $c(z)$ of the flat bundle $\mathcal{H}_{n,m}$. This gives a representation of horizontal sections of the conformal block bundle as hypergeometric integrals. A difficulty here is that a generic condition described in the previous sections does not hold here in general and there is a phenomena of resonance at infinity (see [6] and [20]). We refer the reader to recent work of Looijenga [16] as an interpretation of KZ connections as Gauss–Manin connections and variations of Hodge structures.

Acknowledgment. The author is partially supported by Grant-in-Aid for Scientific Research, Japan Society of Promotion of Science and by World Premier Research Center Initiative, MEXT, Japan.

REFERENCES

- [1] K. Aomoto and M. Kita, *Theory of hypergeometric functions*, Springer Monographs in Mathematics, 2011. DOI: 10.1007/978-4-431-53938-4
- [2] S. Bigelow, Braid groups are linear, *J. Amer. Math. Soc.* **14** (2001), 471 – 486. DOI: 10.1090/S0894-0347-00-00361-1
- [3] S. Bigelow, The Lawrence–Krammer representation, *Topology and geometry of manifolds* (Athens, GA, 2001), 51–68, *Proc. Sympos. Pure Math.* **71** Amer. Math. Soc., 2003.
- [4] E. Date, M. Jimbo, A. Matsuo and T. Miwa, Hypergeometric type integrals and the $sl(2, \mathbf{C})$ Knizhnik–Zamolodchikov equations, *Intern. J. Modern Phys.* **B4** (1990), 1049–1057. DOI: 10.1142/S0217979290000528
- [5] V. G. Drinfel’d, Quasi-Hopf algebras, *Leningrad Math. J.* **1** (1990), 1419–1457.
- [6] B. Feigin, V. Schechtman and A. Varchenko, On algebraic equations satisfied by hypergeometric correlators in WZW models. I, *Comm. Math. Phys.* **163** (1994), 173–184. DOI: 10.1007/BF02101739
- [7] V. G. Knizhnik and A. B. Zamolodchikov, Current algebra and Wess–Zumino models in two dimensions, *Nuclear Phys.* **B247** (1984), 83–103. DOI: 10.1016/0550-3213(84)90374-2
- [8] V. Kac, *Infinite Dimensional Lie Algebras*, Third Edition, Cambridge University Press, 1990. DOI: 10.1017/CBO9780511626234
- [9] T. Kohno, Homology of a local system on the complement of hyperplanes, *Proc. Japan Acad. Ser. A* **62** (1986), 144–147. DOI: 10.3792/pjaa.62.144
- [10] T. Kohno, Monodromy representations of braid groups and Yang–Baxter equations, *Ann. Inst. Fourier* **37** (1987), 139–160. DOI: 10.5802/aif.1114
- [11] T. Kohno, *Conformal Field Theory and Topology*, Translations of Mathematical Monographs, Volume 210 American Mathematical Society, 2002, 182 pages.
- [12] T. Kohno, Hyperplane arrangements, local system homology and iterated integrals, to appear in *Adv. Stud. Pure Math.*

- [13] T. Kohno, Quantum and homological representations of braid groups, to appear in Ann. Scuola Norm. Sup. Pisa.
- [14] D. Krammer, Braid groups are linear, Ann. of Math. **155** (2002), 131–156. DOI: 10.2307/3062152
- [15] R. J. Lawrence, Homological representations of the Hecke algebra, Comm. Math. Phys. **135** (1990), 141 – 191.
- [16] E. Looijenga, A topological interpretation of the KZ system, arXiv:1003.2033.
- [17] I. Marin, Sur les représentations de Krammer génériques, Ann. Inst. Fourier **57** (2007), 1883 – 1925. DOI: 10.5802/aif.2317
- [18] P. Orlik and H. Terao, Arrangements and hypergeometric integrals, MSJ Memoirs **9**, Mathematical Society of Japan, 2001.
- [19] V. Schechtman and A. Varchenko, Hypergeometric solutions of the Knizhnik-Zamolodchikov equation, Lett. in Math. Phys. **20** (1990), 93 – 102. DOI: 10.1007/BF00626523
- [20] R. Silvotti, Local systems on the complement of hyperplanes and fusion rules in conformal field theory, Internat. Math. Res. Notices 1994, no. 3, 111 ff., approx. 17 pp. (electronic).
- [21] A. Varchenko, Determinant formula for Selberg type integrals, Funct. Analysis and Its Appl. **4** (1991), 65 – 66.
- [22] A. Varchenko, Multidimensional hypergeometric functions and representation theory of Lie algebras and quantum groups, Advances in Math. Phys. **21**, World Scientific, 1995.

IPMU, GRADUATE SCHOOL OF MATHEMATICAL SCIENCES, THE UNIVERSITY OF TOKYO, 3-8-1 KOMABA, MEGURO-KU, TOKYO 153-8914 JAPAN

E-mail address: kohno@ms.u-tokyo.ac.jp