

THE MULTIPLICITY POLAR THEOREM, COLLECTIONS OF 1-FORMS AND CHERN NUMBERS

TERENCE GAFFNEY AND NIVALDO G. GRULHA JR.

ABSTRACT. In this work, we show how the Multiplicity Polar Theorem can be used to calculate Chern numbers for collections of 1-forms.

1. INTRODUCTION

Given a space with singularities, and a geometric invariant defined for smooth spaces, it is interesting to see whether or not the invariant is well-defined for the singular space, and, if it is, what are the contributions to the invariant from the singularities.

In a series of papers, Ebeling and Gusein-Zade have discussed the meaning of such invariants as the index of a differential form [10], various notions of the index of a vector field [7], and the Chern numbers, and have described the contributions from the singularities in some cases. In [10], they calculated the radial index of a 1-form on a complete intersection singularity. In [17], it is shown that, in the case of a differential 1-form with an isolated singularity on $X \subset \mathbb{C}^n$, where X is a complex analytic space, that the radial index can be computed using the multiplicity of a pair of modules. The computation of [17] amounts to computing the intersection multiplicity of the graph of the one form ω , which is a subset of the (unprojectivised) conormal bundle of \mathbb{C}^n and the cotangent space of X . In contrast to [10], the calculation is valid for any equidimensional space.

It is clear that this springs from earlier work for vector fields and characteristic classes on singular spaces by Schwartz and Brasselet, ([4, 25]), MacPherson ([24]) Seade and others (see for example [2, 27]). The case of a 1-form is analogous to the case of vector fields, and the indices involved concern the Euler characteristic of the singular variety. This can be regarded as a particular Chern number, and the work of Ebeling and Gusein-Zade for collections of 1-forms extends these notions for other Chern numbers.

In [8], Ebeling and Gusein-Zade developed the notion of the Chern number of a singular space using collections of differential 1-forms. Their numbers are well-defined for any equidimensional reduced complex analytic germ, but they only compute the number for ICIS singularities. Their Chern number is again an intersection number. As in the earlier work, the intersection takes place at the level of conormal spaces; they call the points in X which are the projection of the points of intersection, *special points*.

In the case that we have just one 1-form, the Chern number is the Euler obstruction of the differential form ([7], p. 17). This is related to the Euler obstruction of a set and the Euler obstruction of a function as defined by Brasselet, Massey, Parameswaran and Seade in [3]. In [11], the definition of the Euler obstruction of a function was adapted to the case of 1-form, the Euler obstruction of a function was studied by several authors, in this direction we have for example the papers [21],[27],[6],[17]. In [1], the authors determine relations between the local Euler obstruction of an analytic map f defined in [22] and the Chern number of a convenient collection of 1-forms associated to f .

In this work, we use the multiplicity polar theorem to calculate Chern numbers for any equidimensional reduced complex analytic germ. This extends the earlier work of [17]. For the Chern number problem, one must work with a set of collections of differential 1-forms, and calculate the order of the point where all of the collections are linearly dependent. Since we want to calculate the number of points at which the forms are linearly dependent after a generic

perturbation, this is again a problem involving modules. Because we have a collection of forms we have a collection of modules, so the problem is like a problem in intersection theory, except the spaces are defined by modules not by ideals.

The computation of the Chern numbers is an example of a problem where the underlying vector bundle, which is the tangent bundle in the Chern case, is not defined at every point of X . The set X must be modified; so we pass to the Nash modification of X where the tangent bundle of X is defined in order to understand the problem fully. This process of modifying a space to fill in points where a bundle ξ fails to be defined works in general, and our process of calculating intersection numbers also extends. We outline this in the last section. This suggests the easier problem of calculating Chern numbers when the tangent bundle is well-defined at all points, and more generally, intersection numbers of modules. We take this up in section two. The main themes of section four appear in this material.

Also in Section 2, we recall some basic ideas about the theory of integral closure of modules and the statement of the multiplicity polar theorem.

In Section 3, we recall how Ebeling and Gusein-Zade develop the notion of Chern number in their paper.

In Section 4, we introduce the notion of a special point for a collection. Roughly speaking, a point $p \in X$ is called a special point of the collection $\{\omega_j^{(i)}\}$ of 1-forms on the variety X if there exists a point in the fiber of the Nash modification over p such that the restriction of the 1-forms $\omega_1^{(i)}, \dots, \omega_{d-k_i+1}^{(i)}$ to the point are linearly dependent for each $i = 1, \dots, s$. We next see how special points can be viewed as intersections, and, hence, have an associated intersection number, if isolated. We then begin to solve the “module intersection theoretic problem” for the computation of the Chern numbers described above. We prove a “Gysin” type theorem, (Proposition 4.7) that is, under suitable genericity hypotheses, we can do our calculations on a single space which represents the intersection of all but the last spaces defined by our collection and use the last module associated with the collection restricted to this space for our computations. We also prove a genericity result (Proposition 4.10) which shows that by deforming just the last collection of differential forms, we can ensure the set of collections is generic in an appropriate sense.

We begin Section 5 by recalling a result of Ebeling and Gusein-Zade (Proposition 5.1) relating Chern numbers and special points. In Proposition 5.4 we describe in integral closure terms what it means for $x \in X$ not to be a special point for a collection of forms. After gearing up to apply the multiplicity polar theorem.

In Section 6 we show, in Theorem 6.1, that deforming our last collection allows us to split the contribution of the Chern number from an isolated special point into the multiplicity of a pair of modules and the intersection number of the new collection. Using this as the inductive step, we can write the contribution to the Chern number as a sum of multiplicities of pairs (Corollary 6.2).

We next show that if X is an ICIS, then our formula agrees with that of Ebeling and Gusein-Zade (Corollary 6.3 and the discussion afterwards.)

We close by indicating how our results can be generalized to the case of a bundle E^k defined on a Zariski open, everywhere-dense subset U of an analytic space X , E^k a sub-bundle with k -dimensional fiber of a bundle, F^l , where F^l is defined everywhere.

The authors thank Steven Kleiman for helpful conversations on the connection between their work and the intersection multiplicity of Serre.

2. INTEGRAL CLOSURE OF MODULES

Let (X, x) be a germ of a complex analytic space, X a small representative of the germ, and let \mathcal{O}_X denote the structure sheaf on a complex analytic space X . The study of what it means for a collection of 1-forms to have a special point on a singular space depends on the behavior of limiting tangent hyperplanes. The key tool for studying these limits is the theory of integral closure of modules, which we now introduce.

Definition 2.1. Suppose (X, x) is the germ of a complex analytic space, M a submodule of $\mathcal{O}_{X,x}^p$. Then $h \in \mathcal{O}_{X,x}^p$ is in the integral closure of M , denoted \overline{M} , if for all analytic $\phi : (\mathbb{C}, 0) \rightarrow (X, x)$, $h \circ \phi \in (\phi^*M)\mathcal{O}_1$. If M is a submodule of N and $\overline{M} = \overline{N}$, we say that M is a reduction of N .

To check the definition, it suffices to check along a finite number of curves whose generic point is in the Zariski open subset of X along which M has maximal rank. (Cf. [14])

If a module M has finite colength in $\mathcal{O}_{X,x}^p$, it is possible to attach a number to the module, its Buchsbaum-Rim multiplicity, $e(M, \mathcal{O}_{X,x}^p)$. We can also define the multiplicity $e(M, N)$ of a pair of modules $M \subset N$, M of finite colength in N , as well, even if N does not have finite colength in \mathcal{O}_X^p .

We recall how to construct the multiplicity of a pair of modules using the approach of Kleiman and Thorup [23]. Given a submodule M of a free \mathcal{O}_{X^d} module F of rank p , we can associate a subalgebra $\mathcal{R}(M)$ of the symmetric \mathcal{O}_{X^d} algebra on p generators. This is known as the Rees algebra of M . If (m_1, \dots, m_p) is an element of M , then $\sum m_i T_i$ is the corresponding element of $\mathcal{R}(M)$. Then $\text{Projan}(\mathcal{R}(M))$, the projective analytic spectrum of $\mathcal{R}(M)$, is the closure of the projectivised row spaces of M at points where the rank of a matrix of generators of M is maximal. Denote the projection to X^d by c . If M is a submodule of N or h is a section of N , then h and M generate ideals on $\text{Projan} \mathcal{R}(N)$; denote them by $\rho(h)$ and $\rho(M)$. If we can express h in terms of a set of generators $\{n_i\}$ of N as $\sum g_i n_i$, then in the chart in which $T_1 \neq 0$, we can express a generator of $\rho(h)$ by $\sum g_i T_i / T_1$. Having defined the ideal sheaf $\rho(M)$, we blow it up.

On the blow up $B_{\rho(M)}(\text{Projan} \mathcal{R}(N))$, we have two tautological bundles. One is the pullback of the bundle on $\text{Projan} \mathcal{R}(N)$. The other comes from $\text{Projan} \mathcal{R}(M)$. Denote the corresponding Chern classes by c_M and c_N , and denote the exceptional divisor by $D_{M,N}$. Suppose the generic rank of N (and hence of M) is g .

Then the multiplicity of a pair of modules M, N is:

$$e(M, N) = \sum_{j=0}^{d+g-2} \int D_{M,N} \cdot c_M^{d+g-2-j} \cdot c_N^j.$$

Kleiman and Thorup show that this multiplicity is well-defined at $x \in X$ as long as $\overline{M} = \overline{N}$ on a deleted neighborhood of x . This condition implies that $D_{M,N}$ lies in the fiber over x , hence is compact. Notice that when $N = F$ and M has finite colength in F then $e(M, N)$ is the Buchsbaum-Rim multiplicity $e(M, \mathcal{O}_{X,x}^p)$. There is a fundamental result due to Kleiman and Thorup, the principle of additivity [23], which states that given a sequence of $\mathcal{O}_{X,x}$ -modules $M \subset N \subset P$ such that the multiplicity of the pairs is well-defined, then

$$e(M, P) = e(M, N) + e(N, P).$$

Also if $\overline{M} = \overline{N}$ then $e(M, N) = 0$ and the converse also holds if X is equidimensional. Combining these two results we get that if $\overline{M} = \overline{N}$ then $e(M, N) = e(N, P)$. These results will be used in Section 5.

In studying the geometry of singular spaces, it is natural to study pairs of modules. In dealing with non-isolated singularities, the modules that describe the geometry have non-finite colength, so their multiplicity is not defined. Instead, it is possible to define a decreasing sequence of modules, each with finite colength inside its predecessor, when restricted to a suitable complementary plane. Each pair controls the geometry in a particular codimension.

We also need the notion of the polar varieties of M . The *polar variety of codimension k* of M in X , denoted $\Gamma_k(M)$, is constructed by intersecting $\text{Projan } \mathcal{R}(M)$ with $X \times H_{g+k-1}$ where H_{g+k-1} is a general plane of codimension $g+k-1$, then projecting to X .

Setup: We suppose we have families of modules $M \subset N$, M and N submodules of a free module F of rank p on an equidimensional family of spaces with equidimensional fibers \mathcal{X}^{d+k} , \mathcal{X} a family over a smooth base Y^k . We assume that the generic rank of M, N is $g \leq p$. Let $P(M)$ denote $\text{Projan } \mathcal{R}(M)$, π_M the projection to \mathcal{X} .

We will be interested in computing, as we move from the special point 0 to a generic point, the change in the multiplicity of the pair (M, N) , denoted $\Delta(e(M, N))$. We will assume that the integral closures of M and N agree off a set C of dimension k which is finite over Y , and assume we are working on a sufficiently small neighborhood of the origin, so that every component of C contains the origin in its closure. Then $e(M, N, y)$ is the sum of the multiplicities of the pair at all points in the fiber of C over y , and $\Delta(e(M, N))$ is the change in this number from 0 to a generic value of y . If we have a set S which is finite over Y , then we can project S to Y , and the degree of the branched cover at 0 is $\text{mult}_y S$. (Of course, this is just the number of points in the fiber of S over our generic y .)

Let $C(M)$ denote the locus of points where M is not free, *i.e.*, the points where the rank of M is less than g , $C(\text{Projan } \mathcal{R}(M))$ its inverse image under π_M .

We can now state the Multiplicity Polar Theorem. The proof in the ideal case appears in [15]; the general proof appears in [16].

Theorem 2.2. (*Multiplicity Polar Theorem*) *Suppose in the above setup we have that $\overline{M} = \overline{N}$ off a set C of dimension k which is finite over Y . Suppose further that*

$$C(\text{Projan } \mathcal{R}(M))(0) = C(\text{Projan } \mathcal{R}(M(0))),$$

except possibly at the points which project to $0 \in \mathcal{X}(0)$. Then, for y a generic point of Y ,

$$\Delta(e(M, N)) = \text{mult}_y \Gamma_d(M) - \text{mult}_y \Gamma_d(N),$$

where $\mathcal{X}(0)$ is the fiber over 0 of the family \mathcal{X}^{d+k} , $M(0)$ is the restriction of the module M to $\mathcal{X}(0)$, and $C(\text{Projan } \mathcal{R}(M))(0)$ is the fiber of $C(\text{Projan } \mathcal{R}(M))$ over 0 .

Now, we show how this machinery can be applied to a module intersection problem. Suppose we are given modules $M_1 \subset F_1$ and $M_2 \subset F_2$, F_i free $\mathcal{O}_{X^d, x}$ modules of rank p_i , M_i generated by n_i generators. Suppose $C(M_i)$ is equidimensional, the codimension of $C(M_i)$ is $n_i - p_i + 1$, and the sum of the codimensions is d , $C(M_i)$ equidimensional. If we deform the generators of M_i , how many points do we expect to see where both modules have less than maximal rank?

We can take this number as the intersection number of the two modules.

As further justification, we relate this number to an intersection number at x . Let $\mathcal{M}(p, q)$, $p \leq q$, be the space of $p \times q$ matrices with complex entries and let $D_{p, q}$ be the subspace of $\mathcal{M}(p, q)$ consisting of matrices of rank less than p . The subset $D_{p, q}$ is an irreducible subvariety of $\mathcal{M}(p, q)$ of codimension $q - p + 1$.

Fix a matrix of generators $[M_i]$ of M_i .

Then each matrix $[M_i]$ defines a section Γ_{M_i} of $\mathbb{C}^n \times \mathcal{M}(p_i, n_i)$ in the obvious way; the pair defines a section Γ_{M_1, M_2} of $\mathbb{C}^n \times \mathcal{M}(p_1, n_1) \times \mathcal{M}(p_2, n_2)$. We will assume that

$$X \times D_{p_1, n_1} \times D_{p_2, n_2} \cap \text{Im}(\Gamma_{M_1, M_2})$$

is isolated and lies over x . The intersection number of $X \times D_{p_1, n_1} \times D_{p_2, n_2}$ and $\text{Im}(\Gamma_{M_1, M_2})$ at $(x, \Gamma_{M_1, M_2}(x))$ is the number we want to calculate. In this paper we will abbreviate ‘‘Zariski open set’’ by ‘‘Z-open set’’.

Theorem 2.3. *Suppose each of the sections Γ_{M_i} is transverse to $X^d \times D_{p_i, q_i}$ on a Z-open set U_i such that $U_i \cap C(M_i)$ is Z-open and dense in $C(M_i)$. Then the intersection number of $X \times D_{p_1, n_1} \times D_{p_2, n_2}$ and $\text{Im}(\Gamma_{M_1, M_2})$ at $(x, \Gamma_{M_1, M_2}(x))$ is $e(M_1, \mathcal{O}_{C(M_2), x}) = e(M_2, \mathcal{O}_{C(M_1), x})$.*

Proof. X may be singular, so we assume X is stratified with the canonical Whitney stratification [29]. Then the transversality of Γ_{M_i} means transversality to each $S_j \times D_{p_i, q_i}$, S_j a stratum. This ensures that the generic point of each component of $C(M_i)$ is a smooth point of X . It also ensures that the codimension of $\mathcal{O}_{C(M_i), x}$ is $n_i - p_i + 1$. Since $X \times D_{p_1, n_1} \times D_{p_2, n_2} \cap \text{Im}(\Gamma_{M_1, M_2})$ is isolated and lies over x , the sum of the codimensions of the $C(M_i)$ is d .

Let us show that, at $(x, \Gamma_{M_1, M_2}(x))$, the intersection number of $X \times D_{p_1, n_1} \times D_{p_2, n_2}$ and $\text{Im}(\Gamma_{M_1, M_2})$ is $e(M_1, \mathcal{O}_{C(M_2), x})$. The proof of the other half of the inequality is parallel.

Note that the number of generators of M_1 as a $\mathcal{O}_{C(M_2), x}$ module is

$$n_1 = (n_1 - p_1 + 1) + p_1 - 1 = d - (n_2 - p_2 + 1) + p_1 - 1 = \dim C(M_2) + p_1 - 1.$$

Then, by Theorem 1.2 of [18], we can find a perturbation of $[M_1]$ by a matrix of generic constants such that the section induced by the new matrix, $[\widetilde{M}_1]$, of $C(M_2) \times \mathcal{M}(p_1, n_1)$ is transverse to $C(M_2) \times D_{p, q}$, and the finite number of points at which $[\widetilde{M}_1]$ has less than maximal rank occur at smooth points of $C(M_2)$ and there are $e(M_1, \mathcal{O}_{C(M_2), x})$ of them. In particular, x is no longer a point where both sections have less than maximal rank. (It is not hard to see from the proof of Theorem 2.2 that in fact these lie in the \mathbb{Z} -open dense subset of $C(M_2)$ on which the section Γ_{M_2} is transverse to $X \times D_{p_2, n_2}$.) The transversality conditions on Γ_{M_2} and $\Gamma_{\widetilde{M}_1}$ imply that the section $\Gamma_{\widetilde{M}_1, M_2}$ is transverse to $X \times D_{p_1, n_1} \times D_{p_2, n_2}$ at all points of intersection. The total number of such points counted with multiplicity is the intersection number of $X \times D_{p_1, n_1} \times D_{p_2, n_2}$ and $\text{Im}(\Gamma_{M_1, M_2})$ at $(x, \Gamma_{M_1, M_2}(x))$; the transversality statement implies each point occurs with multiplicity 1. □

Corollary 2.4. Suppose $\mathcal{O}_{X, x}$ is Cohen-Macaulay, then the intersection number of $X \times D_{p_1, n_1} \times D_{p_2, n_2}$ and $\text{Im}(\Gamma_{M_1, M_2})$ at $(x, \Gamma_{M_1, M_2}(x))$ is the colength of the ideal generated by the maximal minors of $[M_i]$, $i = 1, 2$.

Proof. Since $\mathcal{O}_{X, x}$ is Cohen-Macaulay and the structure on $\mathcal{O}_{C(M_2), x}$ given by the minors of $[M_2]$ is generically reduced, it is reduced and $\mathcal{O}_{C(M_2), x}$ is Cohen-Macaulay. Then $e(M_1, \mathcal{O}_{C(M_2), x})$ is the colength of the ideal of minors of $[M_1]$ in $\mathcal{O}_{C(M_2), x}$ which gives the result. □

Looking at the proof of the above theorem, in applying the technique of the proof to geometric problems, we see that we need a description of the desired quantity as an intersection number, and a theorem about the transversality of a deformation of $[M_1]$ by a matrix of generic constants.

If ξ_1 and ξ_2 are vector bundles, we may wish to calculate geometric invariants related to sections of the bundles. If the desired invariant is supported at a point, then locally the sets of sections of our vector bundles are free modules, and we can look at the submodules generated by the given sets of sections. Then the last theorem can be used to calculate the contribution to the invariant at a point where the sections fail to be generic.

In the next couple of sections we will look at a more difficult case, one in which the vector bundle may only be defined on a \mathbb{Z} -open subset of X . This will involve modifying X to produce a new space on which the bundle is defined, then taking into account the fiber of the modification over x .

Before developing these ideas, we mention the connection between the ideas of this section and the intersection multiplicity defined by Serre ([28]). Given modules $M_1 \subset F_1$ and $M_2 \subset F_2$, F_i free $\mathcal{O}_{X^d, x}$ modules of rank p_i as above, Serre's intersection number is the alternating sum of the lengths of the $\text{Tor}^i(F^{p_1}/M_1, F^{p_2}/M_2)$.

Corollary 2.5. Under the hypotheses of Theorem 2.3, Serre's intersection number is the same as $e(M_1, \mathcal{O}_{C(M_2), x}) = e(M_2, \mathcal{O}_{C(M_1), x})$.

Proof. This holds because under small deformations of the M_i the intersection number does not change; but then, by a small deformation, we can reduce to the ideal case (*i.e.*, the modules have rank one less than maximum at common points where they have less than maximal rank). Then, by Theorem 2.3, the intersection number counts the same points as $e(M_1, \mathcal{O}_{C(M_2), x})$. □

For the case where $\mathcal{O}_{X,x}$ is Cohen-Macaulay more can be said. The following result extends some theorems of Buchsbaum and Rim ([5] 2.4 p.207, 4.3 and 4.5 p.223).

Corollary 2.6. Suppose $\mathcal{O}_{X,x}$ is Cohen-Macaulay, then the colength of the ideal generated by the maximal minors of $[M_i]$, $i = 1, 2$, is the length of $F^{p_1}/M_1 \otimes F^{p_2}/M_2$.

Proof. We claim that the complex used to compute the Tor^i is exact, so Serre's intersection number is just the length of $F^{p_1}/M_1 \otimes F^{p_2}/M_2$. To see this, consider the complex for $O_n^{p_1}/M_1$. At points where M_1 has maximal rank, this complex is exact. Further all the maps have maximal rank. These assertions follow because $O_n^{p_1}/M_1 = 0$, and the free resolution of 0 is a trivial complex (lemma 20.1 p491 [13]). Now tensor with $O_n^{p_2}/M_2$, and consider the resulting complex. At points where M_2 has maximal rank we are tensoring with 0, so the complex is exact. At points, different from the origin, where M_2 has less than maximal rank, the complex remains exact, as it is a trivial resolution and the torsion terms are zero as they are independent of the resolution. So the origin is the only point where the complex is not exact; but by the acyclicity lemma, (cf [13] p498) the complex must be exact there as well. The Corollary now follows from the previous one, because since X is Cohen-Macaulay, and $C(M_2)$ has the right dimension, its ring is Cohen-Macaulay as well, so

$$e(M_1, \mathcal{O}_{C(M_2),x})$$

is just the colength of the ideal generated by the maximal minors of $[M_i]$, $i = 1, 2$. \square

3. COLLECTIONS OF 1-FORMS

W. Ebeling and S. M. Gusein-Zade studied indices for collections of 1-forms [7, 8], in this section we will recall some ideas and notation from their papers about these concepts.

If P is a complex analytic manifold of dimension n , then its Euler characteristic $\chi(P)$ is the characteristic number

$$\langle c_n(TP), [P] \rangle = (-1)^n \langle c_n(T^*P), [P] \rangle,$$

where TP is the tangent bundle of the manifold P , T^*P is the dual bundle, and c_n is the corresponding Chern class and $[P]$ the fundamental class of P .

The top Chern class of a vector bundle is the first obstruction to the existence of a non-vanishing section. Other Chern classes are obstructions to the existence of a linearly independent collection of sections. There, instead of 1-forms on a complex variety, we consider collections of 1-forms. Further, to calculate intersections of Chern Classes and hence Chern numbers, we will need collections of collections of 1-forms.

Let $\pi: E \rightarrow P$ be a complex analytic vector bundle of rank m over a complex analytic manifold P of dimension n . It is known that the $(2(n-k))$ -dimensional cycle Poincaré dual to the characteristic class $c_k(E)$ ($k = 1, \dots, m$) is represented by the set of points of the manifold P where $m-k+1$ generic sections of the vector bundle E are linearly dependent.

We continue to use the notation of section two: Let $\mathcal{M}(p, q)$, $p \leq q$, be the space of $p \times q$ matrices with complex entries and let $D_{p,q}$ be the subspace of $\mathcal{M}(p, q)$ consisting of matrices of rank less than p . The subset $D_{p,q}$ is an irreducible subvariety of $\mathcal{M}(p, q)$ of codimension $q-p+1$. The complement $W_{p,q} = \mathcal{M}(p, q) \setminus D_{p,q}$ is the Stiefel manifold of p -frames in \mathbb{C}^q . It is known that the Stiefel manifold $W_{p,q}$ is $2(q-p)$ -connected and $H_{2(q-p)+1}(W_{p,q}) \cong \mathbb{Z}$.

We now develop the notation necessary to handle collections of collections of forms. For the rest of the paper, we will refer to these objects simply as collections.

Let $\mathbf{k} = (k_1, \dots, k_s)$ be a sequence of positive integers with $\sum_{i=1}^s k_i = k$. Consider the space $\mathcal{M}_{m,\mathbf{k}} = \prod_{i=1}^s \mathcal{M}(m-k_i+1, m)$ and the subvariety $D_{m,\mathbf{k}} = \prod_{i=1}^s D_{m-k_i+1, m}$ in it. The variety $D_{m,\mathbf{k}}$ consists of sets $\{A_i\}$ of $(m-k_i+1 \times m)$ matrices such that $\text{rk } A_i < m-k_i+1$ for each $i = 1, \dots, s$. Since $D_{m,\mathbf{k}}$ is irreducible of codimension k , its complement $W_{m,\mathbf{k}} = \mathcal{M}_{m,\mathbf{k}} \setminus D_{m,\mathbf{k}}$ is $(2k-2)$ -connected, $H_{2k-1}(W_{m,\mathbf{k}}) \cong \mathbb{Z}$, and there is a natural choice of a map from an oriented manifold of dimension $2k-1$ to the manifold $W_{m,\mathbf{k}}$.

Let $(X^d, 0) \subset (\mathbb{C}^n, 0)$ be the germ of a purely n -dimensional reduced complex analytic variety at the origin. For $\mathbf{k} = \{k_i\}$, $i = 1, \dots, s$, $j = 1, \dots, d-k_i+1$, let $\{\omega_j^{(i)}\}$ be a collection of germs

of 1-forms on $(\mathbb{C}^n, 0)$. (Note that $\{\omega_j^{(i)}\}$ for a fixed value of i , is itself a collection of $d - k_i + 1$ 1-forms.) Let $\varepsilon > 0$ be small enough so that there is a representative X of the germ $(X, 0)$ and representatives $\{\omega_j^{(i)}\}$ of the germs of 1-forms inside the ball $B_\varepsilon(0) \subset \mathbb{C}^n$.

The kind of points whose multiplicity we wish to compute is described in the next section.

4. SPECIAL POINTS

Definition 4.1. A point $p \in X$ is called a special point of the collection $\{\omega_j^{(i)}\}$ of 1-forms on the variety X if there exists a sequence p_m of points from the non-singular part X_{reg} of the variety X such that the sequence $T_{p_m} X_{\text{reg}}$ of the tangent spaces at the points p_m has a limit L (in $G(d, n)$) and the restriction of the 1-forms $\omega_1^{(i)}, \dots, \omega_{d-k_i+1}^{(i)}$ to the subspace $L \subset T_p \mathbb{C}^n$ are linearly dependent for each $i = 1, \dots, s$. The collection $\{\omega_j^{(i)}\}$ of 1-forms has an isolated special point on $(X, 0)$ if it has no special point on X in a punctured neighborhood of the origin.

Notice that we require each element in the collection to be linearly dependent when restricted to the same limit plane. Notice also, that if an element of the collection has less than maximal rank at a point, then it is linearly dependent on all planes passing through the point.

The framework of this section is a variation on the setting used in [7]. In developing the properties of special points, it is helpful to work on two levels, one of which is based on the Nash modification. The Nash modification comes into play because the tangent bundle of X is not defined at singular points of X . However the Nash bundle is an extension of the tangent bundle on the modified space. We begin to describe this setting.

Let $\{\omega_j^{(i)}\}$ be a collection of germs of 1-forms on $(X, 0)$ with an isolated special point at the origin. Let $\nu : \tilde{X} \rightarrow X$ be the Nash transformation of the variety X , and \tilde{T} the Nash bundle. The collection of 1-forms $\{\omega_j^{(i)}\}$ gives rise to a section $\Gamma(\omega)$ of the bundle

$$\tilde{\mathbb{T}} = \bigoplus_{i=1}^s \bigoplus_{j=1}^{n-k_i+1} \tilde{T}_{i,j}^*$$

where $\tilde{T}_{i,j}^*$ are copies of the dual Nash bundle \tilde{T}^* over the Nash transform \tilde{X} numbered by indices i and j .

Let $\mathbb{D} \subset \tilde{\mathbb{T}}$ be the set of pairs $(x, \{\alpha_j^{(i)}\})$ where $x \in \tilde{X}$ and the collection $\{\alpha_j^{(i)}\}$ is such that $\alpha_1^{(i)}, \dots, \alpha_{n-k_i+1}^{(i)}$ are linearly dependent for each $i = 1, \dots, s$.

Definition 4.2. The local Chern obstruction, $\text{Ch}_{X,0}\{\omega_j^{(i)}\}$, of the collections of germs of 1-forms $\{\omega_j^{(i)}\}$ on $(X, 0)$, at the origin, is the obstruction to extend the section $\Gamma(\omega)$ of the fibre bundle $\tilde{\mathbb{T}} \setminus \mathbb{D} \rightarrow \tilde{X}$ from the preimage of a neighbourhood of the sphere $S_\varepsilon = \partial B_\varepsilon$ to \tilde{X} , more precisely its value, as an element of the cohomology group $H^{2n}(\nu^{-1}(X \cap B_\varepsilon), \nu^{-1}(X \cap S_\varepsilon), \mathbb{Z})$, on the fundamental class of the pair $(\nu^{-1}(X \cap B_\varepsilon), \nu^{-1}(X \cap S_\varepsilon))$.

In the case of a single 1-form, if this is radial, then we are exactly in the setting envisaged by MacPherson to define the local Euler obstruction [24], and otherwise this is essentially the “defect” introduced in [3]. The computation of the local Chern obstruction will be revisited in section 5.

The other setting for the study of special points is closer to X , and we describe it next. This setting will allow us to describe the number of special points as an intersection number.

Let $X^d \subset \mathbb{C}^n$, \mathcal{L}^k be the set of collections of 1-forms respecting the partition of k as above ($k = k_1 + k_2 + \dots + k_s$), $\mathbb{D}_X^k \subset \mathbb{C}^n \times \mathcal{L}^k$ be the closure of the set of pairs $(x, \{l_j^i\})$ such that $x \in X_{\text{reg}}$ and the restriction of the linear functions $l_1^i, \dots, l_{n-k_i+1}^i$ to $T_x X_{\text{reg}} \subset \mathbb{C}^n$ are linearly dependent for each $i = 1, \dots, s$.

Notice that the fiber of \mathbb{D}_X^k over a regular point x of X can be identified with the elements of $M(d - k_i + 1, n)$ which have less than maximal rank when restricted to $T_x X$. Since $T_x X$ is

defined by $n - d$ equations, the fiber of $\mathbb{D}_X^{k_i}$ is itself a fibration over the singular matrices in $M(d - k_i + 1, d)$, hence the restriction of $\mathbb{D}_X^{k_i}$ to the regular points of X has a stratification by rank of the collection restricted to TX_x , $x \in X_{\text{reg}}$.

The collection of 1-forms $\{\omega_j^{(i)}\}$ defines a section Γ_ω of $\mathbb{C}^n \times \mathcal{L}^k$; we will assume in our results that the image of the projection $\pi_X(\mathbb{D}_X^k \cap \text{Im}(\Gamma_\omega))$ is isolated. Note that this implies that the intersection of $\text{Im}(\Gamma_\omega)$ and \mathbb{D}_X^k is isolated as well, since $\pi_X|_{\text{Im}(\Gamma_\omega)}$ is 1-1. We will further assume that the sets $\text{Im}(\Gamma_\omega)$ and \mathbb{D}_X^k have complementary dimension viewed as subsets of $\mathbb{C}^n \times \mathcal{L}^k$. Thus, their intersection number is well-defined.

We are interested in computing this intersection number.

As we will see, this amounts to computing the intersection number of a collection of sets defined by modules. The viewpoint of this paper is to compute this intersection number by successively restricting to the intersection of $k - 1$ elements of the collection. There is a technical condition which describes the way a “good” collection of these sets meet, given in Definition 4.5, which needs some preparation.

Definition 4.3. Given a pair (x, P) , $x \in X$, P in $G(d, n)$, the pair is degenerate for the collection $\{\omega_j\}$, $1 \leq j \leq d - k + 1$, at x , if $\{\omega_j\}|_P$ is linearly dependent at x . Denote the set of degenerate pairs for $\{\omega\}$ by $\mathbb{B}(\omega)$.

Proposition 4.4. *Suppose the collection $\{\omega\}$ is linearly independent at the origin. Then $\mathbb{B}(\omega)$ has codimension k in $X \times G(d, n)$.*

Proof. We can cover $G(d, n)$ with open sets as follows: pick a coordinate plane P of dimension d and a plane of complementary dimension using the complementary coordinates, which we denote by \hat{P} . Clearly, the complementary plane intersects P only at the origin. Consider all planes which are the graphs of a linear map from P to \hat{P} . The equations of these graphs give a unique set of equations describing the plane, and thus associate a matrix of size $(n - d \times n)$ to each plane.

These planes are just the planes that intersect \hat{P} at the origin, and thus are a Zariski open subset of $G(d, n)$.

Suppose U is such an open set, then construct the map from $U \times \text{Hom}(n, d - k + 1)$ to $\text{Hom}(n, n - k + 1)$ by combining the 2 matrices - the element of $\text{Hom}(n, d - k + 1)$ and the matrix of equations describing points of U .

This matrix has size $(n - k + 1 \times n)$ and as a map from

$$U \times \text{Hom}(n, d - k + 1) \rightarrow \text{Hom}(n, n - k + 1)$$

is transverse to the rank stratification. So the codimension of the set of pairs which give matrices of less than maximal rank is $(n) - (n - k + 1) + 1 = k$.

Working globally, it is clear that the set of degenerate pairs is a fibration over the set of elements of $\text{Hom}(n, d - k + 1)$ of maximal rank. So fixing ω we get that the set of degenerate planes has codimension k . (Also fixing a plane P , the set of $L \in \text{Hom}(n, d - k + 1)$ for which the plane is degenerate also has codimension k .) \square

If we have a collection of forms $\{\omega_j^{(i)}\}$ with $\sum_{i=1}^s k_i = k$, every element of which is linearly independent at the origin, then $\mathbb{B}(\omega)$ denotes the planes which are degenerate for every element of the collection. It has codimension less than or equal to k .

Definition 4.5. Given $X^d, 0 \subset \mathbb{C}^n, 0$ with $0 \in S(X)$ and a collection $\{\omega_j^{(i)}\}$ with $\sum_{i=1}^s k_i = k$, $k \leq d$ such that each element of the collection is linearly independent at 0, we say that the collection is proper for X^d if $\dim(\tilde{X}(S(X)) \cap \mathbb{B}(\omega)) \leq d - k - 1$ where $\tilde{X}(S(X))$ is the restriction of the Nash modification of X to $S(X)$, the singular set of X . If this condition holds for a collection of forms linearly independent at 0, with the exception of components of the intersection over the origin, we say the collection is proper on a deleted neighborhood of the origin.

If X is smooth at 0, then we ask $\dim(\tilde{X}(0) \cap \mathbb{B}(\omega)) \leq d - k - 1$.

Remark 4.6. The dimension $\tilde{X}(S(X))$ is at most $d - 1$; if there is a component of dimension $d - 1$, the condition just asks that the component meets $B(\omega)$ properly. Since the dimension of all components of $\tilde{X} \cap B(\omega)$ is at least $d - k$, the properness condition implies that a point of $\tilde{X}(S(X)) \cap \mathbb{B}(\omega)$ is in the closure of points of the intersection lying over smooth points of X . Note also that if $k = d$, and the collection is proper, then $\tilde{X}(S(X)) \cap \mathbb{B}(\omega)$ is empty.

For the geometric description we need of special points, it is necessary to lift our constructions to the level of the Nash modification.

On $\tilde{X} \times \mathcal{L}^k$ we can consider triples (x, P, L) where P is a degenerate plane for L . Call the space of triples \mathbb{D}^k . It is clearly a fibration over \tilde{X} .

Thinking of $\mathbb{C}^n \times G(d, n) \times \mathcal{L}^k$ as a trivial fibration over $\mathbb{C}^n \times G(d, n)$, we have the section induced by ω which we denote by $\Gamma_{\omega, G}$. Note that, if we restrict $\Gamma_{\omega, G}$ to \tilde{X} , then

$$\Gamma_{\omega, G}^{-1}(\mathbb{D}^k) = \mathbb{B}(\omega) \cap \tilde{X}.$$

Now, the image of $\Gamma_{\omega, G}$ has dimension $n + d(n - d)$, while \mathbb{D}^k has codimension $(n - d) + d(n - d) + k$ so the expected dimension of the intersection is $d - k$. Denote the projection of the intersection to X by $S(\omega)$. We can make k a multi-index and make similar constructions; we get the expected dimension of $S(\omega_j^{(i)})$ is $d - (k_1 + \cdots + k_s)$.

Suppose $\{\omega_j^{(i)}\}$ is a collection of 1 forms such that the $\sum k_i = d$ and 0 is an isolated special point. Then all of the various $S(\omega_j^{(i)})$ using different subcollections must have the correct expected dimension; for if $S(\omega_j^{(i)})$ is too large for one subcollection, the excess dimension will be passed to the others and 0 will not be isolated.

Denote $\mathbb{D}^k \cap \text{Im}(\Gamma_{\omega, G})$ by $S_N(\omega)$.

We will also be interested in the notion of a restricted special point; given a collection of 1-forms $\omega_1^{(i)}, \dots, \omega_{d-k_i+1}^{(i)}$, $1 \leq i \leq s$, we say p is a *restricted special point* of the collection if it is a special point, and the sequence of points p_m are in $S(\omega_j^{(i)})$, $1 \leq i \leq s - 1$. In the next proposition we will prove that if the collection $\omega_j^{(i)}$, $1 \leq i \leq s - 1$, is proper, then every special point is a restricted special point.

4.1. Setup. Here we describe our assumption about the collections.

Let $X^d, 0 \subset \mathbb{C}^n, 0$ and $\{\omega_j^{(i)}\}$, a collection of 1-forms with $1 \leq i \leq s$, $1 \leq j \leq d - k_i + 1$, where $\sum k_i = d$.

Assume the collection is arranged so that the first r collections are 1-forms which are linearly independent at 0. We assume the 1-forms in the remaining collection are all linearly dependent at the origin. We assume the collection has an isolated singularity at the origin, and that the generic point of $S(\omega_j^{(i)})$, $1 \leq i \leq s - 1$, is in X_{reg} . If $r = s$ we also assume that the collection made up of the first $s - 1$ elements is proper for X .

Proposition 4.7. *If, in the above set-up, 0 is a isolated special point of the collection, there exists a curve C on $S(\omega_j^{(i)})$, $1 \leq i \leq s - 1$, generically in X_{reg} , such that $\{\omega\}$ is linearly dependent when restricted to the limiting tangent plane T at the origin, and the origin is the only point on $S(\omega_j^{(i)})$ with this property.*

Proof. There are two cases to consider.

Case 1: Assume $r < s$, assume a special point exists. This is also a special point for the collection with the first $s - 1$ elements. Thus $S(\omega_j^{(i)})$, $1 \leq i \leq s - 1$, has positive dimension and its generic point is in X_{reg} . For C , use any curve on $S(\omega_j^{(i)})$, and let T be the limit tangent plane, T_t the tangent plane to X at point t on curve C .

Then $\{\omega_j^{(i)}\}$, $1 \leq i \leq s - 1$, are linearly dependent on T_t , since our point is in $S(\omega_j^{(i)})$; hence, they are linearly dependent on T .

Since $\{\omega_j^{(s)}\}$ is linearly dependent at zero, they are linearly dependent on T also.

Clearly, any point for which such a curve exists is a special point, so the origin is the only such point.

Case 2: $r = s$. Assume we have a special point then $S(\omega_j^{(i)})$, $i \leq s-1$, has positive dimension with generic point in X_{reg} . Denote the limit tangent plane on which all of our collections restrict to be linearly dependent by T .

By the properness assumption, no component of $S_N(\omega)$, $i \leq s-1$, can lie over $S(X)$; for every component of $S_N(\omega)$ must have dimension $d-k$, $k = \sum k_i$, $1 \leq i \leq s-1$, while by the properness assumption the points over $S(X)$ must have dimension $d-1-k$ or less.

This implies that there exists a curve $\varphi: \mathbb{C}, 0 \rightarrow X, 0$ generically in $X_{\text{reg}} \cap S(\omega_j^{(i)})$ with $i \leq s-1$, such that the limiting tangent plane to X along ϕ is T .

Now all the members of our collection are linearly dependent on T including $\{\omega_j^{(s)}\}$. \square

The previous proposition explains why we are interested in collections which are proper. The properness condition means that if we have a special point, then it is a restricted special point as we can realize the limiting plane on which the collection is dependent as a limit of tangents to X along a curve in some $S(\omega_j^{(i)})$. This is the key to our ability to study the intersection number of $\text{Im}(\Gamma_\omega)$ and \mathbb{D}_X^k by restricting to $S(\omega_j^{(i)})$.

There is a converse to the proposition which requires a stronger genericity condition.

Proposition 4.8. *Suppose in the setup of this section, the collection made up of the first $s-1$ elements is proper for X , if the elements of the collection are linearly independent at the origin. If they are not linearly independent, assume they are proper for X on a deleted neighborhood of the origin. Suppose the origin is the only point where there exists a curve C on $S(\omega_j^{(i)})$, $1 \leq i \leq s-1$, generically in X_{reg} , such that $\{\omega\}$ is linearly dependent when restricted to the limiting tangent plane T at the origin. Then the origin is an isolated special point of the collection.*

Proof. Clearly the origin is a special point. If it were non-isolated, then we could apply the previous proposition to find curves detecting the nearby special points as well. \square

These two propositions show that when studying the behavior of special points, with the right genericity requirements, we can restrict from X to $S(\omega_j^{(i)})$, $1 \leq i \leq s-1$, and having done so consider only the last element of the collection.

The next proposition serves as a “moving lemma”.

This proposition and its corollary, together with the multiplicity polar theorem, will show that the invariant of the next section computes the intersection number of $\text{Im}(\Gamma_\omega)$ and \mathbb{D}_X^k . The argument we give is adapted from that appearing in [18], Theorem 1.2, p. 187.

To prove our proposition we want to consider the map:

$$\Theta: \mathbb{D}_X^{k_s} |_{S(\omega^{(i)})_{i \leq s-1}} \times M(d - k_s + 1, n) \rightarrow M(d - k_s + 1, n)$$

given by

$$\Theta((x, L), M) = L - (\omega(x) + M).$$

If we resolve the singularities of the set $\mathbb{D}_X^{k_s} |_{S(\omega)}$, then the composition $\Theta \circ \pi_{\mathbb{D}_X^{k_s}}$ is a submersion because of the contribution from the M term.

In resolving these singularities there may be multiple components. For example, for X^2 a surface in \mathbb{C}^3 with an isolated singularity at the origin, then if $s = 2$ and ω consists of two forms then $\mathbb{D}_X^{k_2} |_{S(\omega)}$ has $(0, M(2, 3))$ as a component. This follows because the polar curve of X^2 is non-empty, which implies that the generic element of $M(2, 3)$ has less than maximal rank on a curve on X^2 , hence lies in $\mathbb{D}_X^{k_2}$ along that curve. However, there will be a unique component for each component of $S(\omega)$ which surjects onto that component. Denote the components of $\mathbb{D}_X^{k_s} |_{S(\omega)}$ which surject onto $S(\omega)$ by $\mathbb{D}_{S(\omega)}^{k_s}$. The fiber of these components over the origin are those collections of forms which are the limits of forms degenerate along a curve in $S(\omega)$.

Let C denote $\Theta^{-1}(0) \cap (\mathbb{D}_{S(\omega)}^{k_s} \times M(d - k_s + 1, n))$ and consider the projection p from C to $M(d - k_s + 1, n)$.

Now

$$\dim C = \dim S(\omega^{(i)})_{i \leq s-1} + \dim(\text{generic fiber of } D^{k_s}) = k_s + [(d - k_s + 1)(n) - k_s].$$

By hypothesis, we have an isolated singularity at 0, so the dimension of $S(\omega^{(i)})_{i \leq s}$ must be 0. This implies the dimension of $S(\omega^{(i)})_{i \leq s-1}$ is k_s , the minimum possible, because otherwise, adding another form to the collection will not lower the dimension of $S(\omega^{(i)})_{i \leq s}$ to 0.

The expression in [] above holds because the codimension of D^{k_s} in $M(\cdot)$ is just

$$-[(d - k_s + 1) + (n - d)] + n + 1 = k_s,$$

so the map $C \rightarrow M(d - k_s + 1, n)$ is a map between equidimensional spaces. Assuming that $0 \in X$ is an isolated special point of the collection, the fiber over 0 of p is a single point $\omega(0)$.

Earlier in this section, we began to look at the intersection number of $\text{Im}(\Gamma_\omega)$ and \mathbb{D}_X^k . Restricting to $S(\omega^{(i)})_{i \leq s-1}$, we can also look at the intersection number of $\text{Im}(\Gamma_{\omega^{(s)}})$ and $\mathbb{D}_{S(\omega)}^{k_s}$. Our moving lemma will be used to calculate this piece of the intersection number of $\text{Im}(\Gamma_\omega)$ and \mathbb{D}_X^k .

Definition 4.9. A special point of a collection $\{\omega_j^{(i)}\}$ of germs of 1-forms on X is non-degenerate, if the section $\Gamma_{\omega_j^{(i)}}, 1 \leq i \leq s$, meets \mathbb{D}_X^k transversally at the point.

We can now state our proposition.

Proposition 4.10. *Given a collection as in the set up of this section, assume that the section $\Gamma_{\omega_j^{(i)}}, 1 \leq i \leq s - 1$, meets \mathbb{D}_X^k transversely on a Z -open subset of $S(\omega^{(i)})_{i \leq s-1}$. Then for generic M , the collection $\{\omega^{(s)} + tM\}$ meets $\mathbb{D}_{S(\omega)}^{k_s}$ transversely at all points close to the origin for t sufficiently small, $t \neq 0$. The number of such points is just the degree of the projection from C to $M(d - k_s + 1, n)$ over the origin in $M(d - k_s + 1, n)$. Further, each such point is a non-degenerate point of the collection $\{\omega_j^{(i)}\}, 1 \leq i \leq s$.*

Proof. Pick M in the complement of the $\Delta(p)$, the discriminant of the projection from C to $M(d - k_s + 1, n)$, such that the line between 0 and M does not intersect these sets close to 0.

Over the points of this line close to 0, the number of points is the degree of p and p is a submersion at each point. This implies that the map obtained by fixing the M term in Θ is a submersion also. Note that the dimension of the source and target of this map are the same, hence the map is in fact a diffeomorphism.

We are interested in exploring the consequences of this fact.

Let us first consider the case where at the points on the fiber of p over tM , t small, x is in the regular part of $S(\omega^{(i)})_{i \leq s-1}$, and the element in \mathbb{D}^{k_s} has rank one less than maximal when restricted to TX_x . Then the resolution of $D^{k_s}|_{S(\omega)}$ is an equivalence at such points because $D^{k_s}|_{S(\omega)}$ is smooth there, so we can work on the tangent space of $D^{k_s}|_{S(\omega)}$. At each point this splits into a direct sum—the part along $S(\omega^{(i)})_{i \leq s-1}$, and the part along the fiber. There is a similar decomposition of the tangent space of the target—the part which can be identified with the fiber in the source, and the normal space to this. The differential is the identity on the tangent space to the fiber, so since the differential is surjective, the restriction of the differential to the tangent space to $S(\omega^{(i)})_{i \leq s-1}$ must surject onto the normal space to the fiber. In turn this implies that the section induced from $\omega + tM$ meets D^k transversely. In fact, since for transversality we just need the tangent vectors to $S(\omega^{(i)})_{i \leq s-1}$, and the other elements of the collection intersect \mathbb{D}_X^k transversely, the collection $\{(\omega^{(i)})_{i \leq s-1}, \omega^s + tM\}$ meets \mathbb{D}_X^k transversely.

In the general case, note that the assumptions we made above coincide with the resolution being an equivalence. If the resolution is not an equivalence, then some tangent vectors on the resolution will be in the kernel of the differential of the projection, hence the differential will not be surjective, contradicting our choice of M . So we only need to consider the above special case.

□

Remark 4.11. If $\{\omega^{(i)}\}$, $1 \leq i \leq r$, is the maximal subset of our collections which meet properly at the origin, then we can choose M so that $\{\omega^{(i)}, \omega^s + tM\}$, $1 \leq i \leq r$, also meet properly at the origin for $t \neq 0$. This will be implicit in our application of our moving lemma.

Denote the collection obtained by moving our last element by $\tilde{\omega}$. From the last proposition we have:

Corollary 4.12. *In the set-up of last proposition we have*

$$\text{Im}(\Gamma_\omega) \cdot \mathbb{D}_X^k = \Gamma(\omega_s) \cdot \mathbb{D}^{k_s}|_{S(\omega^{(i)})_{i \leq s-1}} + \text{Im}(\Gamma_{\tilde{\omega}}) \cdot \mathbb{D}_X^k.$$

Proof. The effect of moving $\{\omega^{(s)}\}$ is to split off points from the intersection $\text{Im}(\Gamma_\omega) \cdot \mathbb{D}_X^k$. The first intersection number on the right is the degree of the projection from C to $M(d - k_s + 1, n)$, and this is the number of points split off from the intersection number on the left hand side of the equation. Moving $\{\omega^{(s)}\}$ ensures that the intersection $\Gamma(\omega_s + tM) \cdot \mathbb{D}^{k_s}|_{S(\omega^{(i)})_{i \leq s-1}}$ is void at the origin, *i.e.*, the intersection point at the origin has split into non-degenerate points. The second term on the right is the remaining points at the origin. □

Corollary 4.13. *In the set-up of last proposition, suppose in addition that the collection $\{\omega^{(i)}\}$, $1 \leq i \leq s - 1$, is proper. Then*

$$\text{Im}(\Gamma_\omega) \cdot \mathbb{D}_X^k = \Gamma(\omega_s) \cdot \mathbb{D}^{k_s}|_{S(\omega^{(i)})_{i \leq s-1}}.$$

Proof. Since the collection $\{\omega^{(i)}, \widetilde{\omega^{(s)}}\}$, $1 \leq i \leq s - 1$, is proper, the intersection of $\Gamma_{\{\omega^{(i)}, \widetilde{\omega^{(s)}}\}}$ and \mathbb{D}_X^k is empty. □

Remark 4.14. If X^{n-1} is a hypersurface and ω_i a collection of forms with $(n - 1) - k + 1$ elements, which are linearly independent at the origin, then it is easy to check if $\tilde{X}(0) \cap B(\omega)$ has dimension $(n - 1) - k - 1$.

Suppose $\dim \tilde{X}(0) \cap B(\omega) \geq (n - 1) - k$. To each point in $B(\omega)$ there corresponds a unique point in $\text{Proj}(\omega)$, the projectivized row space of ω_i .

Note that points of $\text{Proj}(\omega)$ corresponding to points of $\tilde{X}(0) \cap B(\omega)$ are limiting tangent hyperplanes to X at the origin, so the set of points of $\text{Proj}(\omega)$ which are limiting tangent hyperplanes has dimension $\geq (n - 1) - k = \dim \text{Proj}(\omega)$ so every point is a limiting tangent hyperplane.

This is true if and only if $\text{JM}(f, \sum \alpha_i \omega_i)$ fails to be a reduction of $\text{JM}(f) \oplus \mathcal{O}_X$ for all α_i . This can be checked using curves.

Remark 4.15. We continue with the hypersurface isolated singularity case.

Suppose $j + 1$ collections $\{\omega^i\}$ $1 \leq i \leq j + 1$ are in general position *i.e.*, all are linearly independent at 0 and $\dim \cap \text{Proj}(\{\omega^i\})$ is $(n - 1) - \sum_i k_i$. Suppose the properness condition holds for the first j elements but fails for the collection. A dimension count shows that a whole component of $\cap \text{Proj}\{\omega^i\}$ must lie in the fiber of the Nash modification over the origin. Again this is easy to check.

5. COMPUTING CHERN NUMBERS

In this section, we will begin to connect the machinery of section 2 to the computation of Chern numbers of a collection of forms, preparing for the next section which contains our main results.

Ebeling and Gusein-Zade proved this next proposition.

Proposition 5.1. [7] *Let X be a representative of the germ of a complex analytic space, then the local Chern obstruction $\text{Ch}_{X,0}\{\omega_j^{(i)}\}$ of a collection $\{\omega_j^{(i)}\}$ of germs of holomorphic 1-forms is equal to the number of special points on X of a generic deformation of the collection.*

If X is defined by $F : \mathbb{C}^n \rightarrow \mathbb{C}^p$, then the Jacobian module of X denoted $\text{JM}(X)$, is the submodule of \mathcal{O}_X^p generated by the partial derivatives of F . Given a collection of 1-forms with r elements defined on X , form the $p+r$ by n matrix $D(F, \omega)$ by augmenting the Jacobian matrix DF at the bottom with the 1-forms from the collection. Call the submodule of the free module \mathcal{O}_X^{p+r} , generated by the columns of $\begin{pmatrix} D(F) \\ \omega \end{pmatrix}$, the augmented Jacobian module and denote it by $\text{JM}(X, \omega)$.

Note that this construction works in general. Given a submodule M of a free module F , one can select a matrix of generators, and augment the matrix using linear forms. The points at which the new matrix has less than maximal rank is independent of the choice of generators of M as the row space does not change.

In the next lemma, we begin to relate the theory of integral closure and the infinitesimal limiting geometry of our sets of forms.

Lemma 5.2. *Let X be a representative of the germ of a complex analytic space, and let*

$$\mathcal{L} = \{l_1, l_2, \dots, l_s\}$$

be a collection of linear forms. Consider the hyperplanes defined by the forms $\sum a_i l_i$. None of these hyperplanes is a limiting tangent hyperplane to $X, 0$ at the origin if and only if

$$\overline{\text{JM}(X)_p} = \overline{\text{JM}(X)},$$

where p is a submersion whose kernel is the intersection of the kernels of l_1, \dots, l_s . (Here $\text{JM}(X)_p$ is the submodule of $\text{JM}(X)$ generated by $\frac{\partial}{\partial v_i} f$ where the v_i span the kernel of p .)

Proof. Let us prove this result in the special case when p is a linear projection on the last s variables.

If $\text{JM}(X)_p$ is a reduction of $\text{JM}(X)$, then so is $\text{JM}(X)_h$, because $\ker(h) \supset \ker(p)$, where $h = \sum a_i l_i$. Hence, the hyperplane defined by h is not a limiting tangent hyperplane.

Let us prove now that, if $\text{JM}(X)_h = \overline{\text{JM}(X)}$ for all h , then $\overline{\text{JM}(X)_p} = \overline{\text{JM}(X)}$.

Let $K = \ker(p)$, we will show $\text{JM}(X)_p \subset \mathcal{O}_X^k$ is a reduction of $\text{JM}(X) \subset \mathcal{O}_X^k$ if every hyperplane that contains K is not a limiting tangent hyperplane.

Suppose $\text{JM}(X)_p$ is not a reduction. This implies that there exist a map $\phi : (\mathbb{C}, 0) \rightarrow (X, 0)$ and a non-zero $l : \mathbb{C}^n \rightarrow \mathbb{C}$, such that if N is the matrix of generators of $\text{JM}(X)_p$ and M is the matrix of generators of $\text{JM}(X)$, then the ideal $\langle \eta_1(t), \dots, \eta_n(t) \rangle$ generated by the components of $(l \cdot N) \circ \phi(t)$ has larger order than the ideal $\langle m_1(t), \dots, m_n(t) \rangle$ generated by the components of $l \cdot M \circ \phi$. Denote the order of $l \cdot M \circ \phi$ by k . Then

$$\lim 1/t^k \langle m_1(t), \dots, m_n(t) \rangle$$

defines a limiting tangent hyperplane. Since $m_1 = \eta_1, \dots, m_p = \eta_p$, and the order of these terms is greater than k , it follows that T is a limiting tangent hyperplane which contains the kernel of p . □

Given a collection of linear forms $\mathcal{L} = \{l_1, l_2, \dots, l_s\}$, we let $\text{JM}(X, \mathcal{L})$ denote the module whose matrix of generators is gotten by adding as rows $\{l_1, l_2, \dots, l_s\}$ to the jacobian matrix of a set of generators of $I(X)$. In a similar way, let (M, \mathcal{L}) denote the module whose matrix of generators is gotten by adding as rows the $\{l_1, l_2, \dots, l_s\}$ to a matrix of generators of M .

Proposition 5.3. *Let $\mathcal{L} = \{l_1, l_2, \dots, l_s\}$ be a collection of linear 1-forms linearly independent at the origin. Consider the hyperplanes defined by the forms $\sum a_i l_i$. None of these hyperplanes is a limiting tangent hyperplane to $X, 0$ at the origin if and only if $\overline{\text{JM}(X, \mathcal{L})} = \overline{\text{JM}(X)} \oplus \mathcal{O}_X^s$.*

Proof. It suffices to show that $\text{JM}(X, \mathcal{L})$ is a reduction of $\text{JM}(X) \oplus \mathcal{O}_X^s$ if and only if $\text{JM}(X)_p$ is a reduction of $\text{JM}(X)$. Suppose $\text{JM}(X, \mathcal{L})$ is a reduction of $\text{JM}(X) \oplus \mathcal{O}_X^s$, then $\overline{\text{JM}(X, \mathcal{L})}$ contains $\text{JM}(X) \oplus 0$. Restricting to curves, this implies $\overline{\text{JM}(X)_p}$ contains $\text{JM}(X)$.

Suppose $\text{JM}(X)_p$ is a reduction of $\text{JM}(X)$. Then $\overline{\text{JM}(X, \mathcal{L})}$ contains $\text{JM}(X) \oplus 0$. Let $\{v_i\}$ be a collection of vectors such that $l_i(v_j) = \delta_{i,j}$, then $\overline{\text{JM}(X, \mathcal{L})}$ contains $\begin{pmatrix} D(F)(v_i) \\ \mathcal{L}(v_i) \end{pmatrix}$ and $\text{JM}(X) \oplus 0$, so it contains $\text{JM}(X) \oplus \mathcal{O}_X^s$ \square

The previous two propositions can be easily generalized using the same proof. Given M a submodule of a free module F , $\text{Projan } \mathcal{R}(M)$ has a canonical projection to X which is a fibration over the \mathbb{Z} -open subset U_M of X on which M has maximal rank. The fiber of this map consists of hyperplanes. Call the planes in the fibers over U_M , M -planes. The planes in the fibers over $C(M)$ then, are limiting M -planes. Then the analogues of the previous two results are:

Proposition 5.4. *Let X be a representative of the germ of a complex analytic space, and let $\mathcal{L} = \{l_1, l_2, \dots, l_s\}$ be a collection of linear forms linearly independent at the origin. Consider the hyperplanes defined by the forms $\sum a_i l_i$. Then the following statements are equivalent:*

- 1) *None of these hyperplanes is a limiting M -hyperplane to $X, 0$ at the origin.*
- 2) *If p is a submersion whose kernel is the intersection of the kernels of l_1, \dots, l_s then*

$$\overline{M_p} = \overline{M}.$$

- 3) *There is an equality of modules:*

$$\overline{(M, \mathcal{L})} = \overline{M} \oplus \mathcal{O}_X^s.$$

Let $\{\omega_j^{(i)}\}$ be a collection of 1-forms on the variety X , for simplicity we will denote $S(\omega_j^{(i)})$ with $1 \leq i \leq s-1$ by \mathcal{C} . (Recall \mathcal{C} is the set of points where all of the elements of the collection $\{\omega_j^{(i)}\}$ with $1 \leq i \leq s-1$ are singular.) In the next proposition we are interested in characterizing those collections for which the origin is not a special point or restricted special point.

Proposition 5.5. *Let $(X, 0)$ be the germ of an equidimensional reduced analytic variety, X a representative of the germ and $\{\omega_j^{(i)}\}$ a collection of 1-forms; assume the generic point of each component of \mathcal{C} lies in X_{reg} . Assume also the last collection $\{\omega_j^{(s)}\}$ is linearly independent at 0. The origin is not a restricted special point of the collection $\{\omega_j^{(i)}\}$ if and only if $\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}$ is a reduction of $\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$.*

If all of the collections are linearly independent at the origin, and we assume the first $s-1$ elements are proper, then the origin is not a special point of the collection $\{\omega_j^{(i)}\}$ if and only if $\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}$ is a reduction of $\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$.

Proof. Let $\mathcal{L} = \{l_1, l_2, \dots, l_{d-k_s+1}\}$ be a collection of linear 1-forms such that $\omega_i^s(0) = l_i$. As in Lemma 3.3 of [17], using the integral form of Nakayama's lemma we have that $\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}$ is a reduction of $\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$ if and only if $\text{JM}(X, \mathcal{L})|_{\mathcal{C}}$ is a reduction of $\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$. So we can work with \mathcal{L} .

Now we apply the previous proposition, where $M = \text{JM}(X)$ restricted to \mathcal{C} . Then the limiting M -hyperplanes are just the tangent hyperplanes to X as the generic point of each component of \mathcal{C} is in X_{reg} . If some combination of the $\omega_i^s(0) = l_i$ is a limiting tangent hyperplane to X , then that combination is zero when restricted to the limiting tangent plane, and the collection is linearly dependent.

If we assume properness, then since every special point is a restricted special point, the result follows. \square

We will need a refinement of this result for later. The key point in the above argument, is that $\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}$ is a reduction of $\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$ if and only if none of the hyperplanes defined by $\{\omega^{(s)}(0)\}$ is a limiting tangent hyperplane to $X, 0$ at the origin along curves on \mathcal{C} . Given the collection $\{\omega^{(s)}\}$, we can deform it to $\{\omega^{(s)}(0)\}$ by using the linear deformation. This

fixes the one jet of the collection. Denote this family of collections by $\{\omega_L^{(s)}\}$. Denote the family of sections defined by fixing the first $s - 1$ collections and deforming the last one using the linear deformation by $\Gamma_{\{\omega_{s-1}, L\}}$.

Proposition 5.6. *Assume $\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}$ is a reduction of $\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$. Then, the intersection $\text{Im}(\Gamma_{\{\omega_{s-1}, L\}}) \cdot \mathbb{D}_X^k$ is constant in the linear deformation.*

Proof. Since $\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}$ is a reduction of $\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$, the same is true for any member of the family $\{\omega_L^{(s)}\}$. Suppose for some parameter value t_0 that the intersection number changes, i.e., a point splits off. This gives a curve of points in \mathcal{C} , where at each point p , a member of $\{\omega_L^{(s)}\}$ is degenerate when restricted to some plane which is a point over p in the Nash modification. This implies that $\{\omega_L^{(s)}\}(t_0)$ is degenerate when restricted to some plane which is a point over 0 in the Nash modification. As this plane can be reached through a curve on \mathcal{C} , it contradicts that $\text{JM}(X, \omega_L^{(s)}(t_0))|_{\mathcal{C}}$ is a reduction of $\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$. \square

Definition 5.7. $H_{d-1}(X)$, by definition, consists of all elements of \mathcal{O}_X^p which are in the integral closure of $\text{JM}(X^d)$ except at the origin. A related ideal is $H_{c-1}(X, C^c)$ where C is a subset of X of dimension c . It consists of all elements of \mathcal{O}_C^p which are in the integral closure of $\text{JM}(X)$ restricted to C except at the origin.

In general, $H_i(X)$ consists of all elements of \mathcal{O}_X^p which are in the integral closure of $\text{JM}(X)$ off a set of codimension $i + 1$. Sometimes we write $H_i(\text{JM}(X))$. The meaning of $H_i(X, C^c)$ is similar.

Proposition 5.8. *Let $(X, 0)$ be the germ of an equidimensional reduced analytic variety, X a representative of the germ and $\{\omega_j^{(i)}\}$ a collection of 1-forms; assume the generic point of each component of \mathcal{C} lies in X_{reg} . Assume also the last collection $\{\omega_j^{(s)}\}$ is linearly independent at 0. The origin is at most an isolated restricted special point of the collection if and only if $\text{JM}(X, \mathcal{L})|_{\mathcal{C}}$ is a reduction of $H_{c-1}(X, \mathcal{C})|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$ except possibly at x .*

If in addition, the first $s - 1$ collections are proper on a deleted neighborhood of the origin, then the origin is at most an isolated special point.

Proof. Suppose the origin is an isolated restricted special point. Let U be a neighborhood of 0 in X such that x is the only restricted special point. Then by proposition 4.7, $\Gamma_{\omega^{(s)}}$ misses $T^*(X)|_{\mathcal{C}}$ on $U \setminus \{0\}$.

Then $\text{JM}(X, \mathcal{L})|_{\mathcal{C}}$ is a reduction of $\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$ at all $x \in U$, $x \neq 0$ by the previous proposition.

Hence by definition it is a reduction of $H_{c-1}(X, \mathcal{C})|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$ except possibly at x .

On the other hand assume the reduction criterion holds at each point of $U - 0$. This implies $\text{JM}(X, \mathcal{L})|_{\mathcal{C}}$ is a reduction of $\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$ as this last module is a submodule of $H_{c-1}(X, \mathcal{C})|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$. This implies that there are no restricted special points on U except possibly the origin.

If in addition, the first $s - 1$ collections are proper on a deleted neighborhood of the origin, then the lack of restricted special points on $U - 0$ is equivalent to a lack of special points. \square

The last proposition leaves open the question as to whether the origin is a restricted special point if the reduction criterion holds. The next proposition settles this point.

Proposition 5.9. *Suppose the origin is at most an isolated restricted special point. Then*

$$\begin{aligned} & e(\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}, \text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0) \\ &= e(\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}, H_{c-1}(X, \mathcal{C}) \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0) \\ & \quad - e(\text{JM}(X, \mathcal{L})|_{\mathcal{C}}, H_{c-1}(X, \mathcal{C}) \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0) \end{aligned}$$

where \mathcal{L} is a collection of linear 1-forms such that 0 is not a restricted special point for it.

The origin is not a restricted special point if and only if

$$\begin{aligned} & e(\mathrm{JM}(X, \omega^{(s)})|_{\mathcal{C}}, H_{c-1}(X, \mathcal{C}) \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0) \\ &= e(\mathrm{JM}(X, \mathcal{L})|_{\mathcal{C}}, H_{c-1}(X, \mathcal{C}) \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0) \end{aligned}$$

where \mathcal{L} is a collection of linear 1-forms such that 0 is not a restricted special point for it.

Proof. Since the origin is at most an isolated restricted special point all three multiplicities are well-defined. Then, the proof is based on a fundamental result due to Kleiman and Thorup, the principle of additivity [23]. Given a sequence of \mathcal{O}_X modules $A \subset B \subset C$ such that the multiplicity of the pairs is well-defined, then

$$e(A, C) = e(A, B) + e(B, C).$$

The result follows by setting $A = \mathrm{JM}(X, \omega^{(s)})|_{\mathcal{C}}$, $B = \mathrm{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$, and

$$C = H_{d-1}(X, \mathcal{C}) \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}.$$

Using the fact that 0 is not a restricted special point for \mathcal{L} we have that the multiplicity of $(\mathrm{JM}(X, \mathcal{L})|_{\mathcal{C}}, H_{d-1}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1})$ and $(\mathrm{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, H_{d-1}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1})$ are the same.

The origin is not a restricted special point by 5.5 if and only if $\mathrm{JM}(X, \omega^{(s)})|_{\mathcal{C}}$ is a reduction of $\mathrm{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$. The reduction statement holds at 0 if and only if

$$e(\mathrm{JM}(X, \omega^{(s)})|_{\mathcal{C}}, H_{c-1}(X, \mathcal{C}) \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0) = 0,$$

which is true if and only if

$$e(\mathrm{JM}(X, \omega^{(s)})|_{\mathcal{C}}, H_{c-1}(X, \mathcal{C}) \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0) = e(\mathrm{JM}(X, \mathcal{L})|_{\mathcal{C}}, H_{c-1}(X, \mathcal{C}) \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0),$$

where \mathcal{L} is a collection of linear 1-forms such that 0 is not a restricted special point for it. \square

Our next step to apply the Multiplicity Polar Theorem is to show that the polar curve of $\mathrm{JM}(X, \omega^{(s)} + tM)|_{\mathcal{C}}$ is empty.

Proposition 5.10. *Let $X^d, 0 \subset \mathbb{C}^n, 0$ and $\{\omega\}$ a collection of 1-forms $\{\omega_j^{(i)}\}$, $1 \leq i \leq s$, $1 \leq j \leq d - k_i + 1$, $\sum k_i = d$. Assume further the collection has an isolated singularity at the origin, and that the generic point of $S(\omega_j^{(i)})$, $1 \leq i \leq s - 1$ is in X_{reg} . Then, the polar curve of the module $\mathrm{JM}(X, \omega^{(s)} + tM)|_{\mathcal{C}}$ is empty for $\mathfrak{C} = \mathcal{C} \times \mathbb{C}$, where M is a collection of generic linear forms.*

Proof. The polar variety of codimension k of M in X denoted $\Gamma_k(M)$ is constructed by intersecting $\mathrm{Projan} \mathcal{R}(M)$ with $X \times H_{g+k-1}$ where H_{g+k-1} consists of the set of hyperplanes which contain a general plane of dimension $g + k - 1$, and g is the generic rank of $\mathrm{JM}(X, \omega^{(s)} + tM)$, then projecting to X . Note that if M has n generators, so that $\mathrm{Projan} \mathcal{R}(M)$ is contained in $X \times \mathbb{P}^{n-1}$, and the dimension of $\mathrm{Projan} \mathcal{R}(M)$ is greater than or equal to $n + 1$ then the polar varieties of M of codimension n or more are empty, because the codimension of a point in \mathbb{P}^{n-1} is $n - 1$.

With this observation in mind, the next step is to compute the dimension of

$$\mathrm{Projan} \mathcal{R}(\mathrm{JM}(X, \omega^{(s)} + tM)|_{\mathcal{C}}).$$

This dimension is the dimension of the base plus the generic rank of $\begin{pmatrix} D(F) \\ \omega \end{pmatrix}$ minus 1. Now the generic rank of the jacobian matrix is $n - d$, while the generic rank of the jacobian matrix augmented by the $\{\omega_j^s\}$ is $(n - d) + (d - k_s + 1) = n - k_s + 1$. This follows because the generic point of \mathcal{C} is a smooth point of X hence the jacobian matrix has maximal rank there. Because 0 is an isolated singularity, it follows that the augmented matrix generically has maximal rank. Thus we have, since $g = n - k_s + 1$,

$$\dim \mathrm{Projan} \mathcal{R}(\mathrm{JM}(X, \omega^{(s)} + tM)|_{\mathcal{C}}) = k_s + 1 + (n - k_s + 1) - 1 = n + 1.$$

Since the dimension of $\text{Projan } \mathcal{R}(\text{JM}(X, \omega^{(s)} + tM)|_{\mathcal{C}})$ is greater than or equal to the number of generators, there is no polar curve for $\text{JM}(X, \omega^{(s)} + tM)$. \square

Proposition 5.11. *Suppose X is smooth and ω is a 1-form such that ω has a Morse point at 0, then $e(\text{JM}(X, \omega), \text{JM}(X) \oplus \mathcal{O}_X, 0) = 1$.*

Proof. Since X is a smooth manifold, the number of equations of X is $n - d$, so the matrix of generators of $\text{JM}(X, \omega)$ has $n - d + 1$ rows, n columns, and a matrix of generators of $\text{JM}(X) \oplus \mathcal{O}_X$ also has $n - d + 1$ rows with the same $n - d$ first rows. We may assume the equations for X are $z_1 = \cdots = z_{n-d} = 0$.

Then the Rees algebra of $\text{JM}(X) \oplus \mathcal{O}_X$ is $\mathcal{O}_X[S_1, \dots, S_{n-d}, S_{n-d+1}]$, while the ideal corresponding to the inclusion of the Rees algebra of $\text{JM}(X, \omega)$ in that of $\text{JM}(X) \oplus \mathcal{O}_X$ is $(S_i, z_j S_{n-d+1})$ where $1 \leq i \leq n - d$, $n - d < j \leq n$. Now in this example, we know that

$$1 = e(\text{JM}(X, f), \mathcal{O}_X^{n-d+1}) = e(\text{JM}(X, f), \text{JM}(X) \oplus \mathcal{O}_X) + e(\text{JM}(X) \oplus \mathcal{O}_X, \mathcal{O}_X^{n-d+1}),$$

while $e(\text{JM}(X) \oplus \mathcal{O}_X, \mathcal{O}_X^{n-d+1}) = 0$ since the two modules are the same. This uses the additivity of the multiplicity, the fact that ω is Morse on X , and the fact that the multiplicity of $e(\text{JM}(X, \omega), \mathcal{O}_X^{n-d+1})$ is the colength of its ideal of maximal minors.

Now we want to show that we get the same result even if the number of equations is larger than $n - d$. (This happens for example, if we are working at a smooth point of a space which is singular at the origin.) Suppose our choice of generators for $I(X)$ has p generators, $p \geq n - d$. By a change of coordinates we can assume the equations have the form $x_1 = \cdots = x_{n-d} = g_{n-d+1} = \cdots = g_p = 0$, where the matrix of generators of $\text{JM}(X)$ must have the last d columns 0. Then the Rees algebra of $\text{JM}(X) \oplus \mathcal{O}_X$ is the same as before, as is the ideal induced by $\text{JM}(X, \omega)$, so the multiplicity of the pair is the same. \square

Proposition 5.12. *Let $\{\omega_j^{(i)}\}$ be a collection of 1-forms, with $1 \leq i \leq s$, $1 \leq j \leq d - k_i + 1$, $\sum k_i = d$ such that, restricted to X^d , $\{\omega^{(s)}\}$ has a non-degenerate special point at x , x a smooth point of \mathcal{C} and X . Then*

$$e(\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}, \text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}^{n-k_s+1}}, x) = 1.$$

Proof. Let us suppose, that X is a smooth manifold and the number of equations of X is $n - d$ so that the matrix of generators of $\text{JM}(X, \omega^{(s)})$ has $n - k_s + 1$ rows, n columns, and a matrix of generators of $\text{JM}(X) \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}$ also has $n - k_s + 1$ rows with the same $n - d$ first rows. We may assume the equations for X are $x_1 = \cdots = x_{n-d} = 0$. Since we assume \mathcal{C} is smooth at x , and it has dimension k_s , assume that the last k_s coordinates on X define \mathcal{C} . We may assume that the collection $\omega^{(s)}$ has form $\{dx_{n-d+i}, dh\}$ where $1 \leq i \leq d - k_s$ and $h = \sum_{j=1}^{k_s} x_{n-k_s+j}^2$.

As in the last Proposition, in this example, we know that

$$1 = e(\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}, \mathcal{O}_{\mathcal{C}}^{n-k_s+1})$$

$$= e(\text{JM}(X, \omega^{(s)}), \text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}) + e(\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, \mathcal{O}_{\mathcal{C}}^{n-k_s+1}),$$

while $e(\text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, \mathcal{O}_{\mathcal{C}}^{n-k_s+1}) = 0$ since the two modules are the same. This uses the additivity of the multiplicity, the fact that ω is non-degenerate on X , and the fact that the multiplicity of $e(\text{JM}(X, \omega^{(s)}), \mathcal{O}_X^{n-k_s+1})$ is the colength of its ideal of maximal minors, and as in the last Proposition, the general result follows. \square

6. MAIN RESULT

Before giving our main result, it is useful to consider the difference between the case of a vector bundle well-defined at all points, and a bundle like the tangent bundle to a singular space which is not well-defined at $S(X)$. In the second case, we get a special point if $\tilde{X}(0) \cap B(\omega)$ is non-empty. If we alter the last collection of forms, then we can make the last collection generic

on \mathcal{C} , but the singular locus of the modified forms may still be non-empty. In this case the intersection number $\text{Im}(\Gamma_{\tilde{\omega}})$ and \mathbb{D}_X^k may still be non-zero at the origin.

In the first case, the analogue of $\tilde{X}(0)$ consists of a single point, so by altering the last collection of forms we can ensure that the intersection number $\text{Im}(\Gamma_{\tilde{\omega}})$ and \mathbb{D}_X^k is zero at the origin.

This phenomena is the reason that the formula for the Chern numbers for the Nash bundle has many terms, while that of a vector bundle on X has only one.

The next theorem is the key step in the proof of our main result. It allows us to fix each of the collections in turn, until we are left with collections which are linearly independent at the origin and which are proper. Of course, this last collection has no special points.

Theorem 6.1. *Let $(X^d, 0) \subset (\mathbb{C}^n, 0)$ be the germ of an equidimensional reduced analytic variety, with representative X , $\{\omega_j^{(i)}\}$, a collection of 1-forms with $1 \leq i \leq s$, $1 \leq j \leq d - k_i + 1$, $\sum k_i = d$. Assume further the collection has an isolated singularity at the origin, and that the generic point of $S(\omega_j^{(i)})$, $1 \leq i \leq s - 1$, is in X_{reg} . We have that,*

$$\text{Ch}_{X,0}\{\omega_j^{(i)}\} = e(\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}, \text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0) + \text{Im}(\Gamma_{\tilde{\omega}}) \cdot \mathbb{D}_X^k.$$

Proof. Let us consider the family of sets $\mathcal{C} \times \mathbb{C}$. Let $\pi_{\mathcal{C}}$ denote the projection from $\mathcal{C} \times \mathbb{C}$ to \mathcal{C} , and $\pi_{\mathbb{C}}$ the projection to \mathbb{C} . By conservation of number and taking M as in the Proposition 4.10, $\Gamma(\omega^{(s)}) \cdot T^*(X)|_{\mathcal{C}}$ is just

$$\Gamma(\omega^{(s)} + tM) \cdot T^*(X)|_{\mathcal{C}}$$

for t close to 0, and this is just the number of non degenerate special points of

$$\{(\omega^{(s)})_{1 \leq i \leq s-1}, \omega^{(s)} + tM\}$$

for $t \neq 0$, and the intersection number $\text{Im}(\Gamma_{\tilde{\omega}}) \cdot \mathbb{D}_X^k$. (Recall that the collection $\tilde{\omega}$ was defined before Cor 4.12.) To show that the Multiplicity Polar theorem applies, we must also show that

$$C(\text{Projan}(\mathcal{R}(\text{JM}(X \times \mathbb{C}, \omega_t)_{\pi_t}|_{\mathcal{C} \times \mathbb{C}})))(0) = C(\text{Projan}(\mathcal{R}(\text{JM}(X, \omega)|_{\mathcal{C}})))(0),$$

except possibly over $(0, 0) \in \mathcal{C} \times 0$. Since $N = \text{JM}(X)|_{\mathcal{C} \times \mathbb{C}} \oplus \mathcal{O}_{\mathcal{C} \times \mathbb{C}}^{d-k_s+1}$ as a family of modules is independent of t , $\text{Projan } \mathcal{R}(N)$ is a product, hence $C(\text{Projan } \mathcal{R}(N))(0) = C(\text{Projan } \mathcal{R}(N(0)))$. Now, at any point p of $\mathcal{C} \times 0$ close to the origin, there exists a neighborhood U of p such that on U , $\text{JM}(X \times \mathbb{C}, \omega_t)_{\pi_t}|_{\mathcal{C} \times \mathbb{C}} = N$. This implies that over U , $\text{Projan } \mathcal{R}(N)$ is finite over $\text{Projan } \mathcal{R}(\text{JM}(X \times \mathbb{C}, \omega_t)_{\pi_t}|_{\mathcal{C} \times \mathbb{C}})$ and, on $U \cap \mathcal{C} \times 0$, $\text{Projan } \mathcal{R}(N(0))$ is finite over

$$\text{Projan } \mathcal{R}(\text{JM}(X, \omega_0)|_{\mathcal{C}}).$$

Now, since $\text{Projan } \mathcal{R}(\text{JM}(X, \omega_0)|_{\mathcal{C}}) \subset \text{Projan } \mathcal{R}(\text{JM}(X \times \mathbb{C}, \omega_t)_{\pi_t}|_{\mathcal{C} \times \mathbb{C}})(0)$, the desired equality follows, for any element of $\text{Projan } \mathcal{R}(\text{JM}(X \times \mathbb{C}, \omega_t)_{\pi_t}|_{\mathcal{C} \times \mathbb{C}})(0)$ has a preimage in $\text{Projan } \mathcal{R}(N)(0)$ which is $\text{Projan } \mathcal{R}(N(0))$, and the last set maps to $\text{Projan } \mathcal{R}(\text{JM}(X, \omega_0)|_{\mathcal{C}})$. So, the multiplicity polar theorem applies. Note, that since $\text{Projan } \mathcal{R}(N)$ is a product, N has no polar curve, and by 5.10 we know that $\text{JM}(X \times \mathbb{C}, \omega_t)_{\pi_t}|_{\mathcal{C} \times \mathbb{C}}$ has no polar curve either. Now, by Proposition 4.10 we have, $\text{Ch}_{X,0}(\omega_j^{(i)}) = \Gamma(\omega^{(s)} + tM) \cdot T^*(X)|_{\mathcal{C}}$.

Then, using the Multiplicity Polar Theorem we have,

$$\text{Ch}_{X,0}(\omega_j^{(i)}) = e(\text{JM}(X, \omega^{(s)})|_{\mathcal{C}}, \text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0) + \text{Im}(\Gamma_{\tilde{\omega}}) \cdot \mathbb{D}_X^k.$$

□

Suppose the collection is ordered so that the first r collections meet properly and r is the largest integer for which this is true. Let C_i denote $C(\omega^{(1)}, \dots, \omega^{(i)}, \widetilde{\omega^{(s)}}, \dots, \widetilde{\omega^{(i+2)}})$ where $i \leq s - 1$, and $\widetilde{\omega^{(j)}}$ is a collection of generic linear forms so that the collections $\{\omega^{(i)}\}, i \leq r$, $\{\omega^{(j)}\}$ meet properly.

Corollary 6.2. *Suppose the collection is ordered so that the first r collections meet properly and r is the largest integer for which this is true. In the setup of the last Theorem, we have that*

$$\mathrm{Ch}_{X,0}(\omega) = \sum_{i=r}^{i=s-1} e(\mathrm{JM}(X, \omega^{(i+1)})|_{\mathcal{C}_i}, \mathrm{JM}(X)|_{\mathcal{C}_i} \oplus \mathcal{O}_{\mathcal{C}_i}^{d-k_{i+1}+1}, 0)$$

Proof. We prove the Corollary by applying the previous theorem multiple times. First, to $\{\omega^{(s)}|_{\mathcal{C}_{s-1}}\}$, then to $\{\omega^{(s-1)}|_{\mathcal{C}_{s-2}}\}$. Finally, when all but one of our collections meet properly, applying the theorem to $\{\omega^{(r+1)}|_{\mathcal{C}_r}\}$ produces only a single term as the intersection number term is 0. □

Let $C_{i'}$ denote $C(\widetilde{\omega^{(s)}}, \dots, \widetilde{\omega^{(i+2)}})$. Then, $C_{i'}$ is related to the polar varieties of X . For $C(\widetilde{\omega^{(i+2)}})$ is the polar variety of codimension $k(i+2)$, so $C_{i'}$ is the intersection of the corresponding polar varieties. If X is a hypersurface, then in fact this is the polar variety of codimension $\sum_{i+2}^s k(j)$. The hypersurface case is special because since TX_x , $x \in X_0$ has codimension 1, the kernels of all of the $\widetilde{\omega^{(j)}}$ are contained in TX_x if $x \in C_{i'}$, hence x is in the polar variety defined by the union of the kernels.

Corollary 6.3. *In the set up of the last proposition we have*

$$\begin{aligned} \mathrm{Ch}_{X,0}(\omega) &= \sum_{i=r}^{i=s-1} e(\mathrm{JM}(X, \omega^{(i+1)})|_{\mathcal{C}_i}, \mathrm{JM}(X)|_{\mathcal{C}_i} \oplus \mathcal{O}_{\mathcal{C}_i}^{d-k_{i+1}+1}, 0) \\ &= \sum_{i=r}^{i=s-1} e(\mathrm{JM}(X, \omega^{i+1})|_{\mathcal{C}_i}, H_{\mathcal{C}_i-1}(X, \mathcal{C}_i) \oplus \mathcal{O}_{\mathcal{C}_i}^{d-k_{i+1}+1}, 0) \\ &\quad - e(\mathrm{JM}(X, \widetilde{\omega^{i+1}})|_{\mathcal{C}_i}, H_{\mathcal{C}_i-1}(X, \mathcal{C}_i) \oplus \mathcal{O}_{\mathcal{C}_i}^{d-k_{i+1}+1}, 0). \end{aligned}$$

Proof. Apply Proposition 5.9 to expand

$$e(\mathrm{JM}(X, \omega^{(i+1)})|_{\mathcal{C}_i}, \mathrm{JM}(X)|_{\mathcal{C}_i} \oplus \mathcal{O}_{\mathcal{C}_i}^{d-k_{i+1}+1}, 0).$$

□

We want to specialize our results to the case where $X^d, 0 \subset \mathbb{C}^n$ is an ICIS to compare with those of [7]. Given a collection of holomorphic forms ω with an isolated singular point at 0, Ebeling and Gusein-Zade define another notion of index in [7]. In the case $X, 0$ is an ICIS, the index amounts to smoothing X as well as making the forms general, then counting the number of singular points of the new collection on the smoothing. This index is an extension of the GSV-index [20, 28].

This index can be calculated as follows: suppose $\omega_j^{(i)}$, $1 \leq i \leq s$, $1 \leq j \leq n_{k_i} + 1$, $\sum k_i = d$, augment the jacobian matrix of X for each i with $\omega_j^{(i)}$, producing s matrices. Form an ideal in \mathcal{O}_n , using as generators, the generators of $I(X)$, and the maximal minors of the augmented matrices. Denote the resulting ideal by $I_{X, \omega_j^{(i)}}$. Then the index, denoted $\mathrm{ind}_{X,0}(\{\omega\})$ is just the colength of $I_{X, \omega_j^{(i)}}$ in \mathcal{O}_n . ([7], Theorem 20.) Using this index they show that

$$\mathrm{Ch}_{X,0}(\omega_j^{(i)}) = \mathrm{ind}_{X,0}(\{\omega\}) - \mathrm{ind}_{X,0}(\{l\})$$

where $l = \{l_j^{(i)}\}$ is a generic collection of forms. ([7] Cor. 4.)

We will see that this formula can be recovered from the last corollary. If X is an ICIS, and the \mathcal{C}_i have the minimal dimension then the \mathcal{C}_i are Cohen-Macaulay, with ideal the ideal of X and the maximal minors of the augmented matrices. Further, the matrix of generators of $\mathrm{JM}(X)$ has maximal rank except at the origin when restricted to \mathcal{C}_i . This implies that $H_{\mathcal{C}_i-1}(X, \mathcal{C}_i)|_{\mathcal{C}_i}$ is free, so $e(\mathrm{JM}(X, \omega^{(i+1)})|_{\mathcal{C}_i}, H_{\mathcal{C}_i-1}(X, \mathcal{C}_i) \oplus \mathcal{O}_{\mathcal{C}_i}^{d-k_{i+1}+1}, 0) = e(\mathrm{JM}(X, \omega^{(i+1)})|_{\mathcal{C}_i})$. Since $\mathcal{O}_{\mathcal{C}_i}$ is Cohen-Macaulay, the last multiplicity is just the colength of the ideal formed by the maximal

minors of the augmented matrices formed from $\text{JM}(X)$, the collection of forms used to define \mathcal{C}_i and $\omega^{(i+1)}$. It follows that

$$e(\text{JM}(X, \omega^{(i+1)})|_{\mathcal{C}_i}) = \text{ind}(\{\omega^{(1)}, \dots, \omega^{(i)}, \omega^{i+1}, \widetilde{\omega^{(s)}}, \dots, \widetilde{\omega^{(i+2)}}\}).$$

Now the sum on the right hand side of the last corollary telescopes to

$$\text{Ch}_{X,0}(\omega_j^{(i)}) = \text{ind}_{X,0}(\{\omega\}) - \text{ind}(\{\omega^{(1)}, \dots, \omega^{(r)}, \widetilde{\omega^{(s)}}, \dots, \widetilde{\omega^{(r+1)}}\}).$$

Since the last collection on the right hand side is proper, an argument similar to that of Prop. 5.5 shows that the last term is $\text{ind}_{X,0}(\{l\})$.

In the case of surfaces it is not hard to compute with our formula, and we give some examples. As preparation we give two versions of our formula for the case of surfaces which are not ICIS.

The general case of our theorem becomes:

Corollary 6.4. *Let $(X, 0) \subset (\mathbb{C}^n, 0)$ be the germ of a purely 2 dimensional reduced analytic variety, with representative X , $\{\omega\}$, a collection of sets of 1-forms $\{\omega^{(i)}\}$ $1 \leq i \leq 2$, each with two elements. Assume further the collection has an isolated singularity at the origin, and that the generic point of $S(\omega_j^{(1)})$ is in X_{reg} . We have that,*

$$\text{Ch}_{X,0}\{\omega_j^{(i)}\} = e(\text{JM}(X, \omega^{(2)})|_{\mathcal{C}}, \text{JM}(X)|_{\mathcal{C}} \oplus \mathcal{O}_{\mathcal{C}}^{d-k_s+1}, 0) + \text{Im}(\Gamma_{\widetilde{\omega}}) \cdot \mathbb{D}_X^k.$$

Further,

$$\begin{aligned} \text{Ch}_{X,0}\{\omega_j^{(i)}\} &= e(\text{JM}(X, \omega^{(2)})|_{\mathcal{C}_1}, H_0(X, \mathcal{C}_1) \oplus \mathcal{O}_{\mathcal{C}_1}^{n-d}, 0) \\ &- e(\text{JM}(X, \widetilde{\omega^{(2)}})|_{\mathcal{C}_1}, H_0(X, \mathcal{C}_1) \oplus \mathcal{O}_{\mathcal{C}_1}^{n-d}, 0) + e(\text{JM}(X, \omega^{(1)})|_{\mathcal{C}_0}, H_0(X, \mathcal{C}_0) \oplus \mathcal{O}_{\mathcal{C}_0}^{n-d}, 0) \\ &- e(\text{JM}(X, \widetilde{\omega^{(1)}})|_{\mathcal{C}_0}, H_0(X, \mathcal{C}_0) \oplus \mathcal{O}_{\mathcal{C}_0}^{n-d}, 0). \end{aligned}$$

Proof. This is Theorem 5.13 and Corollary 5.15 for the surface case. \square

Further simplification is possible, if X is a complete intersection.

Corollary 6.5. *Suppose in addition X is a complete intersection. Then*

$$\begin{aligned} \text{Ch}_{X,0}\{\omega_j^{(i)}\} &= e(\text{JM}(X, \omega^{(2)})|_{\mathcal{C}_1}, 0) - e(\text{JM}(X, \widetilde{\omega^{(2)}})|_{\mathcal{C}_1}, 0) \\ &+ e(\text{JM}(X, \omega^{(1)})|_{\mathcal{C}_0}, 0) - e(\text{JM}(X, \widetilde{\omega^{(1)}})|_{\mathcal{C}_0}, 0) \\ &= \Gamma^1(\omega^{(1)}) \cdot \Gamma^1(\omega^{(2)}) - \Gamma^1(\widetilde{\omega^{(1)}}) \cdot \Gamma^1(\widetilde{\omega^{(2)}}) \end{aligned}$$

Proof. The first equality holds because X is a complete intersection, and the generic point of \mathcal{C}_i lies in the regular part of X , the Jacobian module of X has maximal rank off the origin, so $H_0(X, \mathcal{C}_1) \oplus \mathcal{O}_{\mathcal{C}_1}^{n-d}$ is just $\mathcal{O}_{\mathcal{C}_1}^n$ so the multiplicity of this pair is just the ordinary Buchsbaum-Rim multiplicity. Since our curves are reduced their rings are Cohen Macaulay, so the multiplicity of $\text{JM}(X, \omega^{(2)})|_{\mathcal{C}_1}$ at 0 is just the colength of the ideal generated by the determinant of the matrix of generators of $\text{JM}(X, \omega^{(2)})|_{\mathcal{C}_1}$. This determinant on X defines the union of $S(X)$ and $\Gamma^1(\omega^{(2)})$ since it does so generically. Thus, this colength is just the intersection of $\Gamma^1(\omega^{(1)})$ with $\Gamma^1(\omega^{(2)})$ and $S(X)$. Applying this insight to each of the terms of the first equality and canceling terms results in the next equality. \square

We give an example using this result.

Example 6.6. *Let $(X, 0) \subset \mathbb{C}^3$ be the germ of a singular surface defined by the function $f : \mathbb{C}^3, 0 \rightarrow \mathbb{C}, 0$, where $f(x, y, z) = y^2 - x^3$. Take the collection of 1-forms $\omega = \{\omega^1, \omega^2\}$, where $\omega^1 = \{(0, x^3, z^2), (z^3, 0, x^2)\}$, and $\omega^2 = \{(y^2, z^3, 0), (0, y^3, z^2)\}$. Then the local Chern obstruction of this collection is 47.*

We will show this using the second equality in the last Corollary. The matrix of generators of $\text{JM}(X, \omega^1)$ and $\text{JM}(X, \omega^2)$ are respectively,

$$\begin{pmatrix} -3x^2 & 2y & 0 \\ 0 & x^3 & z^2 \\ z^3 & 0 & x^2 \end{pmatrix}, \begin{pmatrix} -3x^2 & 2y & 0 \\ y^2 & z^3 & 0 \\ 0 & y^3 & z^2 \end{pmatrix}$$

Calculating the determinants of the matrix of generators of $\text{JM}(X, \omega^2)$ and $\text{JM}(X, \omega^1)$, we get $z^2(2y^3 + 3x^2z^3)$ and $-3x^7 + 2z^5y$. Since we are only interested in the polar curves of ω^i , we use the defining equation to get the equivalent forms $z^2x^2(2xy + 3z^3)$ and $y(-3xy^3 + 2z^5)$. So the equations of the polar curves of our collection are $z^2(2xy + 3z^3) = 0$ and $-3xy^3 + 2z^5 = 0$. To calculate the intersection multiplicity, pull back to the normalization using the map

$$n(t, z) = (t^2, t^3, z).$$

So we want the intersection multiplicity of $z^2(2t^5 + 3z^3) = 0$ and $-3t^{11} + 2z^5 = 0$, which is $(2)(11)+25=47$. Since our underlying space is Whitney equisingular, the polar curves of X are empty, so the term we have computed is the only term in the corollary, so the local Chern obstruction of this collection is 47. (Notice that in this example, one component of the polar of $\text{JM}(X, \omega^2)$ is not reduced. Nonetheless, a careful reading of the proof of our main result shows that in this simple case the main result continues to hold.)

We describe briefly how the work of this section can be generalized. Start with an analytic space X , and a bundle E^k defined on a Zariski open, everywhere-dense subset U of X , E^k a sub-bundle with k -dimensional fiber of a bundle, F^l , where F^l is defined everywhere. Form the relative Nash transformation $N(X, E, F)$ of X as follows: form the bundle over X of k planes in the fiber of F , consider the image of the section of this bundle formed from the fibers of E^k , and take its closure. The relative Nash transformation has a canonical bundle ξ on it which is a sub-bundle of the pullback of F^l to $N(X, E, F)$, ξ and the pullback of E to $N(X, E, F)$ agree restricted to U . By construction and restriction, sections of F^* give sections of $E^*|U$, and ξ^* . If a collection of sections of $E^*|U$ arise in this way from a collection of sections of F^* , and the collection has an isolated special point at $x \in X$, then we can compute the contribution to the Chern number of the dual from our set of sections (hence to ξ) as we did in this section to the dual of the Nash bundle. As in the Nash bundle case, the contribution will be a sum depending on the polar varieties of E relative to F and their intersections. These polar varieties provide some measure of the geometry of E at its singular points on X .

ACKNOWLEDGMENTS

The authors acknowledge the referees for their suggestions and remarks. The second author acknowledges the financial support given by FAPESP grant 2009/08774-0, the program USP-COFECUB grant 07.1.12081.1.7. and CNPq grants 305560/2010-7 and 200430/2011-4.

REFERENCES

- [1] J.-P. Brasselet, N. G. Grulha Jr. and M. A. S. Ruas, *The Euler obstruction and the Chern obstruction*, Bull. Lond. Math. Soc. 42, no. 6, (2010) 1035–1043.
- [2] J.-P. Brasselet, Lê D. T. and J. Seade, *Euler obstruction and indices of vector fields*, Topology 39, no. 6, (2000) 1193–1208. DOI: [10.1016/S0040-9383\(99\)00009-9](https://doi.org/10.1016/S0040-9383(99)00009-9)
- [3] J.-P. Brasselet, D. Massey, A. J. Parameswaran, and J. Seade, *Euler obstruction and defects of functions on singular varieties*, J. London Math. Soc. (2) 70, no. 1, (2004) 59–76.
- [4] J.-P. Brasselet and M.-H. Schwartz, *Sur les classes de Chern d'un ensemble analytique complexe*, Astérisque **82-83** (1981) 93-147.
- [5] D. A. Buchsbaum and D. S. Rim, *A generalized Koszul complex. II. Depth and multiplicity*, Trans. AMS 111 (1963) 197–224.
- [6] J. Fernández de Bobadilla and T. Gaffney *The Lê numbers of the square of a function and their applications* J. London Math. Soc. 77, (2008) 545–557. DOI: [10.1112/jlms/jdm101](https://doi.org/10.1112/jlms/jdm101)
- [7] W. Ebeling and S. M. Gusein-Zade, *Indices of vector fields and 1-forms on singular varieties*, Global aspects of complex geometry, 129169, Springer, Berlin, (2006).

- [8] W. Ebeling and S. M. Gusein-Zade, *Chern obstruction for collections of 1-forms on singular varieties*, Singularity theory, 557–564, World Sci. Publ., Hackensack, NJ, (2007).
- [9] W. Ebeling and S. M. Gusein-Zade, *On the index of a holomorphic 1-form on an isolated complete intersection singularity*, Doklady Math. 64, (2001) 221–224.
- [10] W. Ebeling and S. M. Gusein-Zade, *Indices of 1-forms on an isolated complete intersection singularity*, Moscow Math. J. 3, (2003) 439–455.
- [11] W. Ebeling and S. M. Gusein-Zade, *Radial index and Euler obstruction of a 1-form on a singular variety*, Geom. Dedicata 113, (2005) 231–241. DOI: [10.1007/s10711-005-2184-1](https://doi.org/10.1007/s10711-005-2184-1)
- [12] W. Ebeling, S. M. Gusein-Zade, and J. Seade, *Homological index for 1-forms and a Milnor number for isolated singularities*, Internat. J. Math. 15 (2004) 895–905. DOI: [10.1142/S0129167X04002624](https://doi.org/10.1142/S0129167X04002624)
- [13] D. Eisenbud, *Commutative algebra. With a view toward algebraic geometry*. Graduate Texts in Mathematics, 150. Springer-Verlag, New York, (1995). xvi+785 pp.
- [14] T. Gaffney, *Integral closure of modules and Whitney equisingularity*, Inventiones, 107 (1992) 301–22. DOI: [10.1007/BF01231892](https://doi.org/10.1007/BF01231892)
- [15] T. Gaffney, *Polar methods, invariants of pairs of modules and equisingularity*, Real and Complex Singularities (São Carlos, 2002), Ed. T. Gaffney and M. Ruas, Contemp. Math., 354, Amer. Math. Soc., Providence, RI, June (2004), 113–136.
- [16] T. Gaffney, *The Multiplicity-Polar Theorem*, [arxiv: math/0703650v1](https://arxiv.org/abs/math/0703650v1)
- [17] T. Gaffney, *The Multiplicity Polar Theorem and isolated singularities*, J. Algebraic Geom. 18, no. 3, (2009) 547–574. DOI: [10.1090/S1056-3911-08-00516-X](https://doi.org/10.1090/S1056-3911-08-00516-X)
- [18] T. Gaffney, *The multiplicity of pairs of modules and hypersurface singularities*, Real and Complex Singularities (São Carlos, 2004), Trends in Mathematics, Birkhäuser (2006), 143–168.
- [19] T. Gaffney and S. Kleiman, *Specialization of integral dependence for modules*, Invent. Math. 137 (1999), 541–574. DOI: [10.1007/s002220050335](https://doi.org/10.1007/s002220050335)
- [20] X. Gómez-Mont, J. Seade, and A. Verjovsky, *The index of a holomorphic flow with an isolated singularity*, Math. Ann. **291**, 1991, 737–751. DOI: [10.1007/BF01445237](https://doi.org/10.1007/BF01445237)
- [21] N. G. Grulha Jr., *The Euler Obstruction and Bruce-Roberts’ Milnor Number*, Quart. J. Math. 60(3) (2009) 291–302. DOI: [10.1093/qmath/han011](https://doi.org/10.1093/qmath/han011)
- [22] N. G. Grulha Jr., *L’Obstruction d’Euler Locale d’une Application*, Annales de la Faculté des Sciences de Toulouse, 17, no 1, (2008) 53–71.
- [23] S. Kleiman and A. Thorup, *A geometric theory of the Buchsbaum-Rim multiplicity*, J. Algebra 167 (1994), 168–231. DOI: [10.1006/jabr.1994.1182](https://doi.org/10.1006/jabr.1994.1182)
- [24] R. D. MacPherson, *Chern classes for singular algebraic varieties*, Ann. of Math. 100, (1974), 423–432. DOI: [10.2307/1971080](https://doi.org/10.2307/1971080)
- [25] M.-H. Schwartz, *Champs radiaux sur une stratification analytique*, Travaux en cours, 39, (1991), Hermann, Paris.
- [26] J. Seade, *The index if a vector field on a complex surface with singularities*, in "The Lefschetz Centennial Conf.", ed. A. Verjovsky, Contemp. Math. **58**, Part III, Amer. Math. Soc. 1987, 225–232.
- [27] J. Seade, M. Tibăr and A. Verjovsky, *Milnor Numbers and Euler obstruction*, Bull. Braz. Math. Soc. (N.S.) 36 , no. 2, (2005) 275–283.
- [28] J.P. Serre, *Algèbre Locale. Multiplicities*. Lecture Notes in Mathematics, 11 Springer-Verlag, Berlin-New York (1965).
- [29] B. Teissier, *Variétés polaires locales et conditions de Whitney*, Note aux C. R. Acad. Sc., 290, 1980, 799–802.